

UNITED STATES DEPARTMENT OF THE INTERIOR  
GEOLOGICAL SURVEY

Tabulation of the Ostracode Assemblages and Associated  
Organisms from Selected Bottom Grab Samples Taken in the  
Northeast Gulf of Alaska, F.R.S. Townsend Cromwell Cruise  
EGAL-75KC, 1975: Part II

By

Elisabeth M. Brouwers

Open-File Report 82-949

1982

This report is preliminary and has not been reviewed for  
conformity with U.S. Geological Survey editorial standards.

## CONTENTS

	Page
Introduction.....	1
Methods.....	1
Appendix: Tabulation of Ostracode Assemblages by Sample.....	17
References.....	92

## ILLUSTRATIONS

Figure 1. Locality map of the Gulf of Alaska region studied.....	2
Figure 2. Location map of sample stations between Montague Island and Cape Suckling.....	3
Figure 3. Location map of sample stations between Bering Glacier and Yakutat Bay.....	4

## TABLES

Table 1. List of the selected bottom grab samples by sample type and number.....	5
Table 2. Alphabetical list of the ostracode species occurring in the 126 selected bottom grab samples.....	8
Table 3. Chart showing the presence and absence of the major faunal and floral organisms in the 126 selected bottom grab samples.....	11

## INTRODUCTION

The U.S. Geological Survey is conducting studies of the Alaskan continental shelf, identifying areas and processes of geologic hazard that may affect resource development offshore. These hazards include areas of rapid accumulation of unconsolidated sediment, areas of intense erosional activity, and regions of submarine sliding and slumping; shallow faults are also being identified and classified.

The region of concern in this report is the continental shelf of the northeast Gulf of Alaska, between Montague Island (longitude 148° W.) and Yakutat Bay (longitude 140° W., fig. 1). During 1975, the Fisheries Research Ship Townsend Cromwell collected nearly 350 bottom grab samples from this region.

As part of the northeast Gulf of Alaska project, I am examining the ostracode assemblages (table 2) occurring in the Cromwell bottom grab samples to determine the age, paleoenvironment, and zoogeography. A previous report (Brouwers, 1981) provided a tabulation of the fauna and flora contained in 102 selected bottom grab samples (Van Veen, Shipek and Box Core) from the northeast Gulf of Alaska. The present study similarly records the data from 126 samples collected on the Cromwell cruise (figs. 2, 3; table 1). This information forms a baseline datum of the distribution and relative abundances of the Quaternary ostracode species occurring in the northeast Gulf of Alaska.

## METHODS

The micropaleontological subsamples taken were not standardized, ranging in weight (usually wet) from 200 gm to 1 kg of raw sediment, depending on the initial amount of sample available. All samples were washed on a 200-mesh sieve (75 micrometer openings). The washed sediment was sorted using a set of nested sieves and the sediment was examined for ostracodes 180 micrometers or larger. The tabulation of fauna and flora (tables 2 and 3), therefore, does not include any organism smaller than 180 micrometers in size. All samples were completely "stripped" of valves.

The ostracode adult and juvenile specimens found in each sample were identified at the species level and counted. The percentage each species constitutes of the entire ostracode assemblage from each sample was calculated. The counts refer to the total number of valves or recognizable fragments of a species; a carapace is counted as two valves. Some of the species are represented by specimens containing soft parts, and these are denoted with an asterisk (\*) at the left of a species binomen. These individuals were probably living when collected.

During the microscopic examination and picking of the ostracodes from the sediment, the associated biota was recorded. No attempt was made to determine relative abundance or percentages of these organisms. Table 3 notes their presence or absence in the samples.

The lithologic descriptions included in the appendix were determined on shipboard, as such, should be considered preliminary. This information could reveal faunal associations with particular substrates.



Figure 1.--Locality map of the Gulf of Alaska region studied.




Figure 2.--Location map of sample stations between Montague Island and Cape Suckling.




Figure 3.--Location map of sample stations between Bering Glacier and Yakutat Bay.

Table 1.--List of the selected bottom grab samples by sample type and number  
 [Numbers refer to localities shown in figs. 2 and 3.]

<u>Sample No.</u>	<u>Type</u>	<u>Sample No.</u>	<u>Type</u>
68A	Shipek	104	Shipek
69	Van Veen	105	Box Core
72	Van Veen	106	Van Veen
73	Van Veen	109	Van Veen
74	Van Veen	110	Shipek
75	Van Veen	111	Shipek
76	Van Veen	112	Van Veen
77	Van Veen	113	Shipek
78	Van Veen	115	Shipek
80	Van Veen	117	Shipek
87	Van Veen	118	Shipek
88	Shipek	120	Shipek
89	Shipek	122A	Shipek
91	Van Veen	123	Shipek
94	Van Veen	124A	Box Core
95	Van Veen	124A	Box Core
96	Van Veen	125	Van Veen
97	Van Veen	127	Box Core
98	Van Veen	128	Box Core
99	Van Veen	129	Shipek
103	Shipek	130	Shipek
132	Shipek	164B	Van Veen
133	Shipek	165	Shipek
134	Shipek	166	Shipek

Table 1.--List of the selected bottom grab samples by sample type and number--Continued

<u>Sample No.</u>	<u>Type</u>	<u>Sample No.</u>	<u>Type</u>
138	Shipek	167	Shipek
141	Van Veen	169	Shipek
144 Upper	Van Veen	170B	Box Core
145	Shipek	172	Shipek
146	Shipek	174	Shipek
147	Shipek	175	Shipek
149	Shipek	179B	Box Core
150	Van Veen	181	Shipek
153	Van Veen	203	Van Veen
154	Van Veen	219	Van Veen
155	Van Veen	221	Shipek
157	Van Veen	223	Box Core
158	Van Veen	225	Shipek
159	Shipek	227	Shipek
160A	Van Veen	229	Van Veen
160C	Box Core	231	Van Veen
161	Van Veen	232	Van Veen
162	Van Veen	233	Van Veen
163	Van Veen	237	Van Veen
164A	Van Veen	239	Van Veen
241	Van Veen	405	Shipek
242	Van Veen	406	Shipek
243	Van Veen	407	Shipek
249	Van Veen	409	Shipek
254A	Shipek	412	Shipek

Table 1.--List of the selected bottom grab samples by sample type and number--Continued

<u>Sample No.</u>	<u>Type</u>	<u>Sample No.</u>	<u>Type</u>
258	Van Veen	414	Shipek
264	Shipek	415	Shipek
265	Shipek	422	Shipek
266	Shipek	423	Shipek
268	Shipek	425	Shipek
284	Van Veen	427	Shipek
294	Shipek	428	Shipek
306	Van Veen	429	Shipek
307	Van Veen	431	Shipek
310	Van Veen	432	Shipek
326	Van Veen	433	Shipek
331	Van Veen		
334	Van Veen		
348	Shipek		
349	Shipek		
350	Van Veen		
351	Van Veen		

---

Table 2.--Alphabetical list of the ostracode species recovered from the  
126 selected bottom grab samples

"Acanthocythereis" dunelmensis (Norman, 1865)  
Acuminocythere sp. A  
Ambostacon sp. A  
Argilloecia sp. A  
Argilloecia sp. B  
Aurila sp. A  
"Australicythere" sp. A  
Baffinicythere emarginata (Sars, 1865)  
Bairdia sp. A  
Buntonia sp. A  
Bythocythere sp. A  
Bythocythere sp. B  
Bythocytheromorpha sp. A  
Bythocytheromorpha sp. B  
Candona sp.  
Cluthia cluthae (Brady, Crosskey and Robertson, 1894)  
Cluthia sp. A  
Coquimba sp. A  
Cyclocypris sp.  
Cythere aff. C. alveolivalva Smith, 1952  
Cythere sp. A  
Cytherois sp. A  
Cytheromorpha sp. A  
Cytheromorpha sp. B  
Cytheromorpha sp. D  
Cytheromorpha sp. E  
Cytheropteron aff. C. latissimum of Neale and Howe (1975)  
Cytheropteron aff. C. nodosoalatum Neale and Howe, 1975  
Cytheropteron sp. A  
Cytheropteron sp. B  
Cytheropteron sp. C  
Cytheropteron sp. D  
Cytheropteron sp. E  
Cytheropteron sp. F  
Cytheropteron sp. G  
Cytheropteron sp. H  
Cytheropteron sp. I  
Cytheropteron sp. L  
Cytheropteron sp. M  
Cytheropteron sp. N  
Cytheropteron sp. O  
Cytheropteron sp. Q  
Cytheropteron sp. R  
Cytheropteron sp. S  
Cytheropteron sp. X  
Cytheropteron sp. Y  
Cytheropteron sp. Z  
Cytheropteron sp. AA  
Cytheropteron sp. uniden.  
Cytherura sp. D  
Cytherura sp. F

Table 2.--Alphabetical list of the ostracode species recovered from the  
126 selected bottom grab samples--Continued

Cytherura sp. G  
Cytherura sp. H  
Cytherura sp. I  
Cytherura sp. J  
Elofsonia sp. A  
Eucytherura sp. A  
Finnmarchinella (Barentsovia) angulata (Sars, 1866)  
Finnmarchinella (Barentsovia) sp. A  
Finnmarchinella (Finnmarchinella) finmarchica (Sars, 1866)  
Hemicythere aff. H. quadrinodosa Schornikov, 1974  
Hemicythere sp.  
Hemicytherura sp. A  
Hemicytherura sp. B  
Hemicytherura sp. C  
Krithe sp. A  
"Leguminocythereis" sp. A  
"Leguminocythereis" sp. B  
"Leguminocythereis" sp. D  
Loxoconcha sp. A  
Loxoconcha sp. B  
Loxoconcha sp. D  
Loxoconcha sp. E  
Loxoconcha sp. F  
Macrocypris sp.  
Munseyella sp. A  
Munseyella sp. B  
myodocopid  
Normanicythere sp.  
Palmanella limicola (Norman, 1865)  
Paracypris sp.  
Paradoxostoma aff. P. brunneatum Schornikov, 1975  
Paradoxostoma aff. P. japonicum Schornikov, 1975  
Paradoxostoma sp. B  
Paradoxostoma sp. D  
Paradoxostoma sp. G  
Paradoxostoma sp. H  
Paradoxostoma sp. I  
Paradoxostoma sp. J  
Paradoxostoma sp. uniden.  
Patagonacythere sp. A  
Pectocythere aff. P. parkerae Swain and Gilby, 1974  
Pectocythere aff. P. quadrangulata Hanai, 1957  
Pectocythere sp. D  
Pectocythere sp. E  
Pectocythere sp. F  
Pontocythere sp. A  
Pseudocythere sp.A  
"Radimella" jollaensis (LeRoy, 1943)  
Robertsonites tuberculata (Sars, 1865)  
Schizocythere sp. A  
Sclerochilus sp. C

Table 2.—Alphabetical list of the ostracode species recovered from the  
126 selected bottom grab samples—Continued

Sclerochilus sp. D  
Sclerochilus sp. E  
Sclerochilus sp. F  
Sclerochilus sp. G  
Semicytherura aff. S. undata (Sars, 1965)  
Semicytherura sp. F  
Xestoleberis sp. A  
Xestoleberis sp. B  
Xiphichilus sp.

---

Table 3.--Chart showing the presence and absence of the major faunal and floral

organisms in the 126 selected bottom grab samples

	69	72	73	74	75	76	77	78	80	87	88	89	91	94	95	96	PLANT MATERIAL
DIAATOMS																	X X
FISH PARTS																	
ECHINODERMS																	X
CRUSTACEANS																	
OTHER																	
OSTRACODES																	X X X X X X X X X X X X X X X X
GASTROPODS																	X X X X X X X X X X X X X X X X
PELICYPODS																	X X X X X X X X X X X X X X X X
BRACHIOPODS																	X X X X X X X X X X X X X X X X
WORM TUBES																	X X X X X X X X X X X X X X X X
BRYOZANS																	X X X X X X X X X X X X X X X X
SPICULES																	X X X X X X X X X X X X X X X X
SPONGE																	X X X X X X X X X X X X X X X X
RADIOLARIANS																	X X X X X X X X X X X X X X X X
FORAMINIFERS																	X X X X X X X X X X X X X X X X
PLANCTIC FORAMINIFERS																	X X X X X X X X X X X X X X X X
AGGELLINATE FORAMINIFERS																	X X X X X X X X X X X X X X X X
BENTHIC CALCAREOUS	X																X X X X X X X X X X X X X X X X

Table 3.--Chart showing the presence and absence of the major faunal and floral organisms in the 126 selected bottom grab samples--Continued

	97	98	99	103	104	105	106	109	110	111	112	113	115	117	118	120	122A	123	124B	124A	125
CALCAREOUS																					
BENTHIC																					
FORAMINIFERS																					
AGGLUTINATED																					
BENTHIC																					
FORAMINIFERS																					
PLANKTIC																					
FORAMINIFERS																					
RADOLARIAINS																					
SPONGE																					
SPICULES																					
BRYOZOANS																					
WORM TUBES																					
BRACHIOPODS																					
PELECYPODS																					
GASTROPODS																					
DISTRACODES																					
OTHER CRUSTACEANS																					
ECHINODERMS																					
FISH PARTS																					
DIATOMS																					
PLANT MATERIAL																					

Table 3.--Chart showing the presence and absence of the major faunal and floral organisms in the 126 selected bottom grab samples--Continued

	127	128	129	130	132	133	134	138	141	144U	145	146	147	149	150	153	154	155	157	158
CALCAREOUS BENTHIC FORAMINIFERS																				
AGGLOMERATED BENTHIC FORAMINIFERS																				
BENTHIC FORAMINIFERS																				
PLANKTIC FORAMINIFERS																				
SPONGE SPICULES																				
RADIOLARIANS																				
BRYOZANS																				
WORM TUBES																				
BRACHIOPODS																				
PELLETICPODS																				
GASTROPODS																				
OSTRACODES																				
OTHER CRUSTACEANS																				
ECHINODERMS																				
FISH PARTS																				
DIAATOMS																				
PLANT MATERIAL																				

Table 3.--Chart showing the presence and absence of the major faunal and floral

organisms in the 126 selected bottom grab samples--Continued

	159	160 A	160 C	161	162	163	164 A	164 B	165	166	167	169	170B	172	174	175	179B	181	203	219	221	223	225
MATERIAL																							
PLANT																							
DIAATOMS				x																			
FISH PARTS																	x						
ECHINODERMS								x	x	x	x	x	x										
CRUSTACEANS			x																				
OTHER			x																				
OSTRACODES		x					x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
GASTROPODS							x			x	x	x	x	x	x	x	x	x	x	x	x	x	x
PELICYPODS							x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
BRACHIOPODS																							
WORM TUBES			x	x	x											x							
BRYOZANS		x															x	x		x			
SPONGE																	x	x					
RADIOLARIANS																	x						
FORAMINIFERS			x																				
PLANKTIC																							
FORAMINIFERS																							
BENTHIC																							
AGGLOMINATED																	x						
FORAMINIFERS																							
BENTHIC																							
CALCAREOUS	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	

Table 3.—Chart showing the presence and absence of the major faunal and floral

organisms of the 126 selected bottom grab samples--Continued

	227	229	231	232	233	237	239	241	242	243	249	254A	258	264	265	266	268	284	294	306	307	310	326	331	334
CALCAREOUS BENTHIC FORAMINIFERS	X			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
AGGLOMINTATED BENTHIC FORAMINIFERS																									
PLANKTIC FORAMINIFERS																									
PLANKTIC FORAMINIFERS																									
RADIOLARIANS																									
SPONGE																									
SPICULES																									
BRYOZANS																									
WORM TUBES																									
BRACHIOPODS																									
PELICYPODS																									
GASTROPODS																									
OSTRACODES																									
OTHER CRUSTACEANS																									
ECHINODERMS																									
FISH PARTS																									
DIAATOMS																									
PLANT MATERIAL																									

Table 3.—Chart showing the presence and absence of the major faunal and floral

organisms of the 126 selected bottom grab samples--Continued

## APPENDIX

EGAL-75-KC

Shipek-68A

Latitude: 59° 42.6' N  
 Longitude: 146° 15.0' W  
 Water Depth: 81 meters

Lithology: Gray mud with gravel and shells.

Organisms: Calcareous Benthic Foraminifers  
 Sponge Spicules  
 Cheiostome Bryozoans  
 Brachiopods  
 Pelecypods  
 Ostracodes

Ostracode Species:	Adult	Juv.	%
" <u>Australicythere</u> " sp. A	39	136	51.17
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	30	11	11.99
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	22	11	9.65
<u>Pectocythere</u> aff. <u>P. parkerae</u> Swain and Gilby, 1974	19	4	6.73
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	3	13	4.68
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)		9	2.63
<u>Cytheromorpha</u> sp. B	4	1	1.46
<u>Ambostracon</u> sp. A	1	3	1.17
<u>Cytheropteron</u> sp. G	4		1.17
<u>Cytheropteron</u> sp. L	2	2	1.17
<u>Cytheropteron</u> sp. H	3	1	1.17
<u>Cytheropteron</u> sp. E	2	2	1.17
<u>Eucytherura</u> sp. A	3		0.88
<u>Argilloecia</u> sp. A	3		0.88
<u>Hemicytherura</u> sp. B	2		0.59
<u>Cytheropteron</u> sp. N	2		0.59
<u>Finmarchinella</u> ( <u>Barentosovia</u> ) sp. A	2		0.59
<u>Cytheropteron</u> sp. F	1	1	0.59
<u>Finmarchinella</u> ( <u>Finmarchinella</u> ) <u>finmarchica</u> (Sars, 1866)	1		0.29
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	1		0.29
<u>Acuminocythere</u> sp. A	1		0.29
<u>Cytheropteron</u> sp. D	1		0.29
<u>Sclerochilus</u> sp. C		1	0.29
<u>Sclerochilus</u> sp. F	1		0.29
Total Ostracodes	342		

EGAL-75-KC

Van Veen-69

Latitude: 60° 16.6' N  
 Longitude: 146° 14.6' W  
 Water Depth: 49 meters

Lithology: Olive-gray, sandy, clayey silt.

Organisms: Ostracodes

Ostracode Species:	Adults	Juv.	%
* <u>"Leguminocythereis"</u> sp. A	20	198	20.20
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	150	48	18.35
<u>Pectocythere</u> sp. D	159	38	18.26
<u>Loxoconcha</u> sp. A	59	88	13.62
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)	15	54	6.39
<u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)	7	49	5.19
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	19	29	4.45
<u>Cytheropteron</u> sp. A	18	23	3.40
* <u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974	4	27	2.87
<u>Cytheromorpha</u> sp. B	15	2	1.58
<u>Cytheromorpha</u> sp. E	16		1.48
<u>Cytheromorpha</u> sp. A	14		1.30
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952	2	7	0.83
<u>Candona</u> sp.	1	5	0.56
<u>Elofsonia</u> sp. A	4		0.37
<u>Cytherois</u> sp. B	2		0.19
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe (1975)	1		0.09
<u>Cythere</u> sp. A		1	0.09
<u>Cyclocypris</u> sp.	1		0.09
<u>Acuminocythere</u> sp. A		1	0.09
<u>Cytherura</u> sp. J	1		0.09
<u>Cytheropteron</u> sp. I		1	0.09
Total Ostracodes	1079		

EGAL-75-KC

Van Veen-72

Latitude: 60° 15.3' N  
 Longitude: 146° 00.8' W  
 Water Depth: 90 meters

Lithology: Gray-brown mud.

Organisms:      Calcareous Benthic Foraminifers  
                  Pelecypods  
                  Gastropods  
                  Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Robertsonites tuberculata</u> (Sars, 1865)	5	61	33.67
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	1	52	27.04
<u>Cytheropteron</u> sp. B	8	14	11.23
<u>Bythocythere</u> sp. B	8	12	10.20
<u>Cytheropteron</u> sp. A	3	11	7.14
<u>Palmanella limicola</u> (Norman, 1865)	1	6	3.57
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	4	3	3.57
<u>Cytheromorpha</u> sp. B		2	1.02
<u>Cytherois</u> sp. A	1	1	1.02
<u>Cyclocypris</u> sp.		1	0.51
<u>Pseudocythere</u> sp. A		1	0.51
<u>Loxoconcha</u> sp. B	1		0.51
Total Ostracodes	196		

EGAL-75-KC      Van Veen-73

Latitude:    60° 10.45' N  
 Longitude:   146° 01.35' W  
 Water Depth: 95 meters

Lithology:      Gray-brown, clayey silt.

Organisms:      Calcareous Benthic Foraminifers  
                  Agglutinated Benthic Foraminifers  
                  Planktic Foraminifers  
                  Pelecypods  
                  Ostracodes  
                  Diatoms

Ostracode Species:	Adults	Juv.	%
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	24	155	72.47
<u>Palmanella limicola</u> (Norman, 1865)	14	5	7.69
<u>Loxoconcha</u> sp. A	8	6	5.67
<u>Eucytherura</u> sp. A	7	1	3.24

Ostracode Species:	Adult	Juv.	%
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975		7	2.83
<u>Munseyella</u> sp. A	5		2.02
<u>Cluthia</u> <u>cluthae</u> (Brady, Crosskey, and Robertson, 1874)	4		1.62
<u>Cytheropteron</u> sp. D	3	1	1.62
<u>Cytheropteron</u> sp. Q		3	1.22
<u>Loxoconcha</u> sp. B	2		0.81
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)		1	0.41
" <u>Australicythere</u> " sp. A		1	0.41
Total Ostracodes	247		

EGAL-75-KC                  Van Veen-74

Latitude:    60° 09.2' N  
 Longitude:   146° 01.5' W  
 Water Depth: 90 meters

Lithology:      Gray-brown silt.

Organisms:        Calcareous Benthic Foraminifers  
                     Proteinaceous Worm Tubes  
                     Gastropods  
                     Ostracodes  
                     Diatoms

Ostracode Species:	Adults	Juv.	%
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	2	23	32.47
<u>Loxoconcha</u> sp. A	5	14	24.68
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	9	9	23.38
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975	5	3	10.39
<u>Eucytherura</u> sp. A	3		3.90
<u>Cytheropteron</u> sp. L		1	1.30
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)	1		1.30
<u>Cytheropteron</u> sp. D	1		1.30
<u>Munseyella</u> sp. B	1		1.30
Total Ostracodes	77		

EGAL-75-KC

Van Veen-75

Latitude: 60° 07.4' N  
Longitude: 146° 02.3' W  
Water Depth: 84 meters

Lithology: Rounded cobbles at the surface, underlain by gray mud with pea-sized gravel.

Organisms: Calcareous Benthic Foraminifers  
Planktic Foraminifers  
Sponge Spicules  
Pelecypods  
Ostracodes  
Echinoderm Spines  
Diatoms

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)	1	2	42.86
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	1	2	42.86
<u>Palmanella limicola</u> (Norman, 1865)		1	14.29
Total Ostracodes 7			

EGAL-75-KC

Van Veen-76

Latitude: 60° 02.0' N  
Longitude: 146° 00.5' W  
Water Depth: 77 meters

Lithology: 10 cm of gravel and shell fragments overlying gray silty mud.

Organisms: Calcareous Benthic Foraminifers  
Sponge Spicules  
Proteinaceous and Agglutinated Worm Tubes  
Brachiopods  
Pelecypods  
Gastropods  
Pteropod  
Ostracodes  
Echinoderm Fragments

Ostracode Species:	Adults	Juv.	%
<u>"Australicythere"</u> sp. A	25	111	51.52
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	12	20	12.12
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1857	8	15	5.68
<u>Pectocythere</u> aff. <u>P. parkerae</u> Swain and Gilby, 1974	11	4	5.68
<u>Cytheropteron</u> sp. G	11	4	5.68
<u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)	2	3	1.89
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)	2	3	1.89
<u>Cytheromorpha</u> sp. B	4	1	1.89
<u>Cytheropteron</u> sp. H	1	4	1.89
<u>Eucytherura</u> sp. A		3	1.14
<u>Cytheropteron</u> sp. N	1	2	1.14
<u>Hemicytherura</u> sp. B	2		0.76
<u>Munseyella</u> sp. A	2		0.76
<u>Ambostracon</u> sp. A		2	0.76
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975		1	0.38
<u>Hemicytherura</u> sp. A	1		0.38
<u>Cytheropteron</u> sp. F		1	0.38
<u>"Leguminocythereis"</u> sp. D		1	0.38

Total Ostracodes 264

EGAL-75-KC Van Veen-77

Latitude: 59° 56' N  
 Longitude: 146° 1.5' W  
 Water Depth: 86 meters

Lithology: Gray, pebbly mud.

Organisms: Calcareous Benthic Foraminifers  
 Agglutinated Benthic Foraminifers  
 Planktic Foraminifers  
 Sponge Spicules  
 Pelecypods  
 Gastropods  
 Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	5	4	25.71
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	3	6	25.71
<u>Cytheropteron</u> sp. H	1	2	8.57
<u>Cytheropteron</u> sp. E	1	2	8.57
<u>Munseyella</u> sp. A	2		5.71
<u>Eucytherura</u> sp. A		2	5.71
<u>Cytheropteron</u> sp. G		2	5.71
<u>Palmanella limicola</u> (Norman, 1865)	1		2.86
<u>"Australicythere"</u> sp. A		1	2.86
<u>Cytheropteron</u> sp. O	1		2.86
<u>Cytheromorpha</u> sp. A	1		2.86
<u>Cytheropteron</u> sp. F		1	2.86
Total Ostracodes 35			

EGAL-75-KC                  Van Veen-78

Latitude: 59° 51.6' N  
Longitude: 146° 00.9' W  
Water Depth: 101 meters

Lithology: Gray mud.

Organisms:                  Calcareous Benthic Foraminifers  
                                 Agglutinated Benthic Foraminifers  
                                 Proteinaceous Worm Tubes  
                                 Ostracodes  
                                 Echinoderm Fragments  
                                 Diatoms

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)		38	67.86
<u>Palmanella limicola</u> (Norman, 1865)	5	5	17.86
<u>Cytherois</u> sp. A	2		3.57
<u>Eucytherura</u> sp. A	1	1	3.57
<u>Cytheropteron</u> sp. Q	1		1.79
<u>Cytheropteron</u> sp. D	1		1.79
<u>Munseyella</u> sp. B	1		1.79
<u>Cytheropteron</u> sp. A		1	1.79
Total Ostracodes 56			

EGAL-75-KC

Van Veen-80

Latitude: 59° 46.7' N  
 Longitude: 145° 59.5' W  
 Water Depth: 91 meters

Lithology: Tan-gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Agglutinated Benthic Foraminifers  
 Sponge Spicules  
 Proteinaceous Worm Tubes  
 Pelecypods  
 Gastropods  
 Ostracodes  
 Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Palmanella limicola</u> (Norman, 1865)	12	2	18.18
<u>Cytheropteron aff. C. latissimum</u> of Neale and Howe (1975)	3	10	16.88
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	1	11	15.58
<u>"Australicythere" sp. A</u>		6	7.79
<u>Pectocythere aff. P. parkerae</u> Swain and Gilby, 1974	2	3	6.49
<u>Cluthia</u> sp. A	3	1	5.20
<u>Cytheropteron</u> sp. H		3	3.90
<u>Cytheropteron</u> sp. F		2	2.60
<u>Robertsonites tuberculata</u> (Sars, 1865)		2	2.60
<u>Hemicytherura</u> sp. A		2	2.60
<u>Paradoxostoma</u> sp. G	2		2.60
<u>Cytheromorpha</u> sp. A	2		2.60
<u>Eucytherura</u> sp. A		1	1.30
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975		1	1.30
<u>Cytheropteron</u> sp. O		1	1.30
<u>Cytheropteron</u> sp. E		1	1.30
<u>Cytheropteron</u> sp. D	1		1.30
<u>Paradoxostoma</u> sp.		1	1.30
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	1		1.30
<u>Aurila</u> sp. A		1	1.30
<u>Sclerochilus</u> sp. C		1	1.30
<u>Cytheropteron</u> sp. G	1		1.30

Total Ostracodes 77

EGAL-75-KC

Van Veen-87

Latitude: 60° 06.9' N  
Longitude: 145° 34.4' W  
Water Depth: 126 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Planktic Foraminifers  
Radiolarians  
Proteinaceous Worm Tubes  
Pelecypod Fragments  
Ostracodes  
Diatoms

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	25		52.08
<u>Robertsonites tuberculata</u> (Sars, 1865)	11		22.92
<u>Krithe</u> sp. A	1	1	4.17
<u>Munseyella</u> sp. A	2		4.17
<u>Cytheropteron</u> sp. Q	2		4.17
<u>Loxoconcha</u> sp. B	2		4.17
<u>Cytheropteron</u> sp. R	1		2.08
<u>Palmanella limicola</u> (Norman, 1865)	1		2.08
<u>Bythocythere</u> sp. B	1		2.08
<u>Cluthia</u> sp. A	1		2.08
Total Ostracodes 48			

EGAL-75-KC

Shipek-88

Latitude: 59° 59.2' N  
Longitude: 145° 34.0' W  
Water Depth: 88 meters

Lithology: Gray mud with pebbles and cobbles.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Sponge Spicules  
Proteinaceous Worm Tubes  
Pelecypods  
Pteropod  
Ostracodes  
Echinoderm Fragments

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	4	5	50.0
<u>Munseyella</u> sp. A	2		11.11
" <u>Australicythere</u> " sp. A		1	5.56
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)		1	5.56
<u>Cytheropteron</u> sp. E	1		5.56
<u>Eucytherura</u> sp. A	1		5.56
<u>Pectocythere</u> aff. <u>P. parkerae</u> Swain and Gilby, 1974	1		5.56
<u>Cytheropteron</u> sp. N		1	5.56
<u>Cytheropteron</u> sp. F		1	5.56
Total Ostracodes 18			

EGAL-75-KC                    Shipek-89

Latitude:        59° 58.5' N  
 Longitude:      145° 34.2' W  
 Water Depth:    84 meters

Lithology:       Gray mud with cobbles and gravel.

Organisms:       Calcareous Benthic Foraminifers  
                   Sponge Spicules  
                   Proteinaceous Worm Tubes  
                   Cheiostome Bryozoans  
                   Pelecypods  
                   Gastropods  
                   Pteropod  
                   Ostracodes  
                   Echinoderm Fragments

Ostracode Species:	Adults	Juv.	%
<u>Cluthia</u> <u>cluthae</u> (Brady, Crosskey, and Robertson, 1874)	2		100
Total Ostracodes 2			

EGAL-75-KC

Van Veen-91

Latitude: 59° 50.5' N  
Longitude: 145° 39.6' W  
Water Depth: 97 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Planktic Foraminifers  
Proteinaceous Worm Tubes  
Ostracodes  
Echinoderm Fragments  
Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Palmanella limicola</u> (Norman, 1865)	5	1	50.00
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)		3	25.00
<u>Munseyella</u> sp. A	1		8.33
<u>Eucytherura</u> sp. A		1	8.33
<u>Munseyella</u> sp. B		1	8.33
Total Ostracodes	12		

EGAL-75-KC

Van Veen-94

Latitude: 60° 07.7' N  
Longitude: 145° 21.0' W  
Water Depth: 97 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Proteinaceous Worm Tubes  
Pelecypods  
Gastropods  
Ostracodes  
Organic Debris

Ostracode Species:	Adults	Juv.	%
* <u>Robertsonites tuberculata</u> (Sars, 1865)		24	42.11
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	2	18	35.09
<u>Loxoconcha</u> sp. A	1	3	7.02
<u>Cluthia</u> sp. A	3		5.26
<u>Palmanella limicola</u> (Norman, 1865)	1		1.75
<u>Loxoconcha</u> sp. B	1		1.75
<u>Bythocythere</u> sp. B	1		1.75
<u>Cytheropteron</u> sp. B	1		1.75
<u>Cytheropteron</u> sp. Q		1	1.75
<u>Pseudocythere</u> sp. A		1	1.75

Total Ostracodes 57

EGAL-75-KC Van Veen-95

Latitude: 60° 03.3' N  
Longitude: 145° 19.8' W  
Water Depth: 132 meters

Lithology: Greenish-gray mud.

Organisms: Calcareous Benthic Foraminifers  
Proteinaceous Worm Tubes  
Pelecypods  
Pteropod  
Ostracodes

Ostracode Species:	Adults	Juv.	%
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	3	21	80.00
<u>Palmanella limicola</u> (Norman, 1865)	1	1	6.67
* <u>Krithe</u> sp. A	1		3.33
<u>Cytheropteron</u> sp. B	1		3.33
<u>Robertsonites tuberculata</u> (Sars, 1865)		1	3.33
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975		1	3.33

Total Ostracodes 30

EGAL-75-KC

Van Veen-96

Latitude: 59° 59.2' N  
Longitude: 145° 19.3' W  
Water Depth: 119 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Proteinaceous Worm Tubes  
Pelecypods  
Ostracodes  
Echinoderm Fragments  
Diatoms

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	1	8	56.25
<u>Palmanella limicola</u> (Norman, 1865)	1	1	12.50
<u>Krithe</u> sp. A	2		12.50
<u>Robertsonites tuberculata</u> (Sars, 1865)		1	6.26
<u>Eucytherura</u> sp. A		1	6.25
<u>Loxoconcha</u> sp. B		1	6.25
Total Ostracodes 16			

EGAL-75-KC

Van Veen-97

Lithology: 59° 55.7' N  
Longitude: 145° 19.5' W  
Water Depth: 101 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Proteinaceous Worm Tubes  
Pelecypods  
Gastropods  
Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Loxoconcha</u> sp. A	4	2	28.57
<u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)	2	3	23.81
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	5		23.81
<u>Eucytherura</u> sp. A	1	1	9.52
<u>Cytheropteron</u> sp. L		1	4.76
<u>Munseyella</u> sp. A	1		4.76
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975		1	4.76
Total Ostracodes 21			

EGAL-75-KC Van Veen-98

Latitude: 59° 52.5' N  
 Longitude: 145° 19.8' W  
 Water Depth: 101 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Agglutinated Benthic Foraminifers  
 Proteinaceous Worm Tubes  
 Pelecypods  
 Ostracodes  
 Echinoderm Fragments  
 Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	7	2	30.00
<u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)	1	6	23.33
<u>Eucytherura</u> sp. A	4		13.33
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)	1	2	10.0
<u>Loxoconcha</u> sp. A	1	1	6.67
<u>Cytheropteron</u> sp. D	2		6.67
<u>Munseyella</u> sp. A	2		6.67
<u>Cluthia</u> <u>cluthae</u> (Brady, Crosskey, and Robertson, 1874)	1		3.33
Total Ostracodes 30			

EGAL-75-KC              Van Veen-99

Latitude:    59° 50.4' N  
Longitude:   145° 20.6' W  
Water Depth: 110 meters

Lithology:      Gray mud.

Organisms:      Calcareous Benthic Foraminifers  
                    Agglutinated Benthic Foraminifers  
                    Proteinaceous Worm Tubes  
                    Pelecypods  
                    Pteropods  
                    Ostracodes  
                    Echinoderm Fragments  
                    Diatoms

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)	5	18	57.50
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	5	6	27.50
* <u>Munseyella</u> sp. A	4		10.00
* <u>Cluthia</u> sp. A	1		2.50
<u>Xestoleberis</u> sp. B		1	2.50
Total Ostracodes 40			

EGAL-75-KC              Shipek-103

Latitude:    60° 09.4' N  
Longitude:   144° 58.2' W  
Water Depth: 35 meters

Lithology:      Thin layer of gray mud overlying dark, fine-grained sand.

Organisms:      Calcareous Benthic Foraminifers  
                    Proteinaceous Worm Tubes  
                    Pelecypods  
                    Gastropods  
                    Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>"Leguminocythereis"</u> sp. A	3	2	41.67
* <u>Loxoconcha</u> sp. A	4		33.33
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)	2		16.67
<u>Pectocythere</u> sp. D	1		8.33
Total Ostracodes 12			

EGAL-75-KC

Shipek-104

Latitude: 60° 08.1' N  
Longitude: 144° 54.9' W  
Water Depth: 53 meters

Lithology: Gray mud with some sand lenses.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Planktic Foraminifers  
Proteinaceous Worm Tubes  
Pelecypods  
Gastropod  
Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	49	98	34.83
* <u>Robertsonites tuberculata</u> (Sars, 1865)	31	107	32.70
<u>Pectocythere aff. P. quadrangulata</u> Hanai, 1957	81	22	24.41
<u>Palmanella limicola</u> (Norman, 1865)	13	6	4.50
<u>Loxoconcha</u> sp. A	11	3	3.32
<u>Cytheropteron</u> sp. A		1	0.24

Total Ostracodes 422

EGAL-75-KC

Box Core-105

Latitude: 59° 57.1' N  
Longitude: 144° 55.4' W  
Water Depth: 183 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Planktic Foraminifers  
Pelecypods  
Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	6	63	38.33
* <u>Loxoconcha</u> sp. B	52	2	30.00
* <u>Cytheropteron</u> sp. A	6	12	10.00
* <u>Cluthia</u> sp. A	13		7.22
* <u>Krithe</u> sp. A	8	4	6.67
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	2	1	1.67
<u>Cytheropteron</u> sp. M	3		1.67
<u>Cytheropteron</u> sp. Q	1	1	1.11
<u>Bythocythere</u> sp. B	2		1.11
<u>Cytherura</u> sp. I	1		0.56
<u>"Leguminocythereis"</u> sp. A		1	0.56
<u>Cluthia</u> <u>cluthae</u> (Brady, Crosskey, and Robertson, 1874)	1		0.56
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)		1	0.56

Total Ostracodes 180

EGAL-75-KC Van Veen-106

Latitude: 59° 57.0' N  
Longitude: 144° 57.4' W  
Water Depth: 192 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Pelecypods  
Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	3	47	31.25
* <u>Loxoconcha</u> sp. B	44	2	28.75
* <u>Cytheropteron</u> sp. A	12	13	15.63
<u>Bythocythere</u> sp. B	4	9	8.13
* <u>Krithe</u> sp. A	11	2	8.13
<u>Cytheropteron</u> sp. Q	2	5	4.38
<u>Cluthia</u> sp. A	5		3.13
<u>Cytheropteron</u> sp. AA	1		0.63

Total Ostracodes 160

EGAL-75-KC

Van Veen-109

Latitude: 59° 43.4' N  
 Longitude: 144° 52.7' W  
 Water Depth: 102 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Agglutinated Benthic Foraminifers  
 Planktic Foraminifers  
 Proteinaceous Worm Tubes  
 Bryozoans  
 Pelecypods  
 Ostracodes  
 Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Palmanella limicola</u> (Norman, 1865)	1	20	17.65
<u>Pectocythere</u> aff. <u>P. parkerae</u> Swain and Gilby, 1974	1	13	11.77
<u>Loxoconcha</u> sp. A	3	7	8.40
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952		10	8.40
<u>Acuminocythere</u> sp. A		8	6.72
" <u>Leguminocythereis</u> " sp. A		7	5.88
<u>Cytheromorpha</u> sp. B	3	4	5.88
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	1	5	5.04
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)		6	5.04
<u>Cytheromorpha</u> sp. A	4	2	5.04
<u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974		4	3.36
<u>Buntonia</u> sp. A	2	1	2.52
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	1	2	2.52
<u>Loxoconcha</u> sp. D		2	1.68
<u>Cytheropteron</u> sp. E		2	1.68
<u>Cytheromorpha</u> sp. E	2		1.68
<u>Cytheropteron</u> sp. A		2	1.68
<u>Cythere</u> sp. A		1	0.84
* <u>Sclerochilus</u> sp. F	1		0.84
" <u>Australicythere</u> " sp. A		1	0.84
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975	1		0.84
<u>Ambostracon</u> sp. A		1	0.84
<u>Cytherois</u> sp. A		1	0.84
Total Ostracodes 119			

EGAL-75-KC

Shipek-110

Latitude: 59° 41.5' N  
Longitude: 144° 47.2' W  
Water Depth: 97 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Ostracodes  
Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Robertsonites tuberculata</u> (Sars, 1865)	1	5	22.22
<u>Cytheropteron</u> sp. G	4		14.82
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	3	1	14.82
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	2	1	11.11
<u>Loxoconcha</u> sp. A		2	7.41
<u>Munseyella</u> sp. A	2		7.41
<u>Cytheropteron</u> sp. F	1		3.70
<u>Palmanella limicola</u> (Norman, 1865)		1	3.70
<u>Cytheromorpha</u> sp. A	1		3.70
<u>Cytheropteron</u> sp. N	1		3.70
<u>Sclerochilus</u> sp. C		1	3.70
<u>Paradoxostoma</u> sp.		1	3.70
Total Ostracodes 27			

EGAL-75-KC

Shipek-111

Latitude: 59° 38.9' N  
Longitude: 144° 41.0' W  
Water Depth: 148 meters

Lithology: Olive mud overlying gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Planktic Foraminifers  
Proteinaceous Worm Tubes  
Brachiopods  
Pelecypods  
Gastropods  
Ostracodes  
Echinoderm Fragments

Ostracode Species:	Adults	Juv.	%
<u>Palmanella limicola</u> (Norman, 1865)	5	6	24.44
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	2	7	20.00
<u>Robertsonites tuberculata</u> (Sars, 1865)		8	17.78
<u>Eucytherura</u> sp. A	2	2	8.89
<u>Cytheropteron</u> sp. L	2	2	8.89
<u>Munseyella</u> sp. A	3		6.67
<u>Cytheropteron</u> sp. D	1		2.22
<u>Cluthia</u> sp. A	1		2.22
<u>Buntonia</u> sp. A		1	2.22
" <u>Leguminocythereis</u> " sp. A		1	2.22
<u>Munseyella</u> sp. B	1		2.22
<u>Loxoconcha</u> sp. B	1		2.22

Total Ostracodes 45

EGAL-75-KC Van Veen-112

Latitude: 59° 37.6' N  
 Longitude: 144° 37.0' W  
 Water Depth: 145 meters

Lithology: Olive-green mud overlying gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Agglutinated Benthic Foraminifers  
 Pelecypods  
 Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Munseyella</u> sp. A	3		23.08
* <u>Palmanella limicola</u> (Norman, 1865)	2	1	23.08
* <u>Macrocypris</u> sp.	2		15.39
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)		2	15.39
<u>Robertsonites tuberculata</u> (Sars, 1865)		1	7.69
<u>Cytheropteron</u> sp. L	1		7.69
<u>Eucytherura</u> sp. A	1		7.69

Total Ostracodes 13

EGAL-75-KC

Shipek-113

Latitude: 59° 34.9' N  
Longitude: 144° 30.0' W  
Water Depth: 139 meters

Lithology: Olive mud with occasional small pebbles and coarse sand overlying gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Planktic Foraminifers  
Radiolarians  
Sponge Spicules  
Calcareous and Proteinaceous Worm Tubes  
Pelecypods  
Gastropod  
Ostracodes  
Echinoderm Fragments

Ostracode Species:	Adults	Juv.	%
<u>Sclerochilus</u> sp. D	5		55.56
<u>Krithe</u> sp. A	3		33.33
<u>Eucytherura</u> sp. A	1		11.11
Total Ostracodes 9			

EGAL-75-KC

Shipek-115

Latitude: 59° 46.0' N  
Longitude: 144° 47.7' W  
Water Depth: 64 meters

Lithology: Dark gray, muddy, fine-grained sand.

Organisms: Calcareous Benthic Foraminifers  
Worm Tubes  
Bryozoans  
Pelecypods  
Ostracodes  
Echinoderm Fragments

Ostracode Species:	Adults	Juv.	%
<u>Loxoconcha</u> sp. A	40	93	15.13
* <u>"Leguminocythereis"</u> sp. A	9	104	12.86
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	67	39	12.06
<u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974	3	89	10.47
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952	7	48	6.26
* <u>Cytheromorpha</u> sp. B	39	6	5.12
* <u>Palmanella limicola</u> (Norman, 1865)	15	23	4.32
<u>Aurila</u> sp. A	3	34	4.21
<u>Pectocythere</u> sp. D	9	25	3.87
<u>Robertsonites tuberculata</u> (Sars, 1865)	3	28	3.53
<u>Pectocythere</u> aff. <u>P. parkerae</u> Swain and Gilby, 1974	3	22	2.84
" <u>Radimella</u> " <u>jollaensis</u> (LeRoy, 1943)		18	2.05
<u>Acuminocythere</u> sp. A	3	14	1.93
<u>Cytheromorpha</u> sp. E	15		1.71
<u>Cythere</u> sp. A	2	11	1.48
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975	1	11	1.37
<u>Paradoxostoma</u> sp. G	8	4	1.37
<u>Cytheropteron</u> sp. A	7	5	1.37
* <u>Buntonia</u> sp. A	7	4	1.25
<u>Hemicytherura</u> sp. A	10	1	1.25
<u>Loxoconcha</u> sp. D	7	2	1.02
<u>Coquimba</u> sp. A	2	6	0.91
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	1	5	0.68
<u>Cytherois</u> sp. A	5		0.57
* <u>Paradoxostoma</u> sp. I	4	1	0.57
<u>Hemicytherura</u> sp. B	5		0.57
<u>Ambostracon</u> sp. A		4	0.46
<u>Cytheropteron</u> sp. Z	1	2	0.34
* <u>Argilloecia</u> sp. A	2		0.23
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975) myodocopid		1	0.11
Total Ostracodes 879	1		0.11

EGAL-75-KC

Shipek-117

Latitude: 59° 43.0' N  
 Longitude: 144° 38.1' W  
 Water Depth: 119 meters

Lithology: Gray mud.

Organisms:      Calcareous Benthic Foraminifers  
                   Agglutinated Benthic Foraminifers  
                   Planktic Foraminifers  
                   Pelecypods  
                   Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Palmanella limicola</u> (Norman, 1865)	2	7	33.33
* <u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	4	4	29.63
<u>Pectocythere</u> sp. D	1	2	11.11
<u>Robertsonites tuberculata</u> (Sars, 1865)		2	7.41
<u>"Leguminocythereis"</u> sp. A		2	7.41
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	1		3.70
<u>Loxoconcha</u> sp. A		1	3.70
<u>Cytheropteron</u> sp. S	1		3.70
Total Ostracodes 27			

EGAL-75-KC      Shipek-118

Latitude:      59° 40.7' N  
 Longitude":     144° 33.3' W  
 Water Depth:    137 meters

Lithology:      Gray mud.

Organisms:      Calcareous Benthic Foraminifers  
                   Agglutinated Benthic Foraminifers  
                   Planktic Foraminifers  
                   Radiolarians  
                   Proteinaceous Worm Tubes  
                   Pelecypods  
                   Ostracodes  
                   Occasional Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Eucytherura</u> sp. A	5	2	36.84
<u>Palmanella limicola</u> (Norman, 1865)	3	2	26.32
<u>Robertsonites tuberculata</u> (Sars, 1865)		2	10.53
<u>Cytheropteron</u> sp. L		1	5.26
<u>Cytheropteron</u> sp. D	1		5.26
<u>Munseyella</u> sp. B	1		5.26
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)		1	5.26

Ostracode Species:	Adult	Juv.	%
<u>"Leguminocythereis"</u> sp. A		1	5.26
Total Ostracodes 19			

EGAL-75-KC              Shipek-120

Latitude:    59° 48.8' N  
 Longitude:   144° 41.0' W  
 Water Depth: 66 meters

Lithology:      Dark, fine-grained sand.

Organisms:        Calcareous Benthic Foraminifers  
                   Agglutinated and Proteinaceous Worm Tubes  
                   Pelecypods  
                   Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>"Leguminocythereis"</u> sp. A	159	109	78.82
<u>Pectocythere</u> sp. D	30		8.82
* <u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	14	1	4.41
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)	5	4	2.65
<u>"Leguminocythereis"</u> sp. B	2	3	1.87
<u>Loxoconcha</u> sp. A	3		0.88
<u>Eucythere</u> sp. A	2		0.59
<u>Cytheromorpha</u> sp. B	2		0.59
<u>Buntonia</u> sp. A		2	0.59
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952		1	0.29
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	1		0.29
<u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)		1	0.29
<u>Aurila</u> sp. A		1	0.29
Total Ostracodes 340			

EGAL-75-KC

Shipek-122A

Latitude: 59° 55.6' N  
Longitude: 144° 31.4' W  
Water Depth: 55 meters

Lithology: Gray mud.

Organisms:      Calcareous Benthic Foraminifers  
                  Agglutinated Benthic Foraminifers  
                  Proteinaceous Worm Tubes  
                  Pelecypods  
                  Ostracodes  
                  Echinoderm Fragments  
                  Plant Material  
                  Diatoms

Ostracode Species:	Adults	Juv.	%
* <u>Palmanella limicola</u> (Norman, 1865)	52	21	51.41
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	2	18	14.09
* <u>Buntonia</u> sp. A	11	6	11.97
<u>Pontocythere</u> sp. A		5	3.52
<u>Cluthia</u> sp. A	4		2.82
<u>Cytheropteron</u> sp. D	4		2.82
<u>Hemicythere</u> aff.. <u>quadrinodosa</u> Schornikov, 1974		4	2.82
<u>Cythere</u> sp. A		4	2.82
<u>Cythere</u> aff. C. <u>alveolivalva</u> Smith, 1952		3	2.11
<u>Cytheromorpha</u> sp. B		2	1.41
<u>Cluthia</u> <u>cluthae</u> (Brady, Crosskey, and Robertson, 1874)	1		0.70
" <u>Leguminocythereis</u> " sp. A		1	0.70
<u>Cytherois</u> sp. A		1	0.70
<u>Loxoconcha</u> sp. A	1		0.70
<u>Cytheropteron</u> sp. Y		1	0.70
<u>Lamelliscapha</u> sp.	1		0.70

Total Octaneade 143

EGAL-75-KC

Shipek-123

Latitude: 59° 56.7' N  
Longitude: 144° 40.2' W  
Water Depth: 210 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Radiolarians  
Gastropod  
Ostracodes

Ostracode Species:	Adults	Juv.	%
* <u>Loxoconcha</u> sp. B	11	2	30.23
<u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)	9		20.93
<u>Cytheropteron</u> sp. A	4	3	16.28
<u>Cluthia</u> sp. A	4		9.30
<u>Cytheropteron</u> sp. Q	2	2	9.30
<u>Bythocythere</u> sp. B	2	1	6.98
<u>Cytheropteron</u> sp. C	2		4.65
<u>"Leguminocythereis"</u> sp. A		1	2.33
Total Ostracodes 43			

EGAL-75-KC

Box Core-124B

Latitude: 59° 57.5' N  
Longitude: 144° 43.2' W  
Water Depth: 234 meters

Lithology: Sticky, firm, gray mud with some fragments of more firm mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Ostracodes  
Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron</u> sp. Q	1		100
Total Ostracodes 1			

EGAL-75-KC

Box Core-124A

Latitude: 59° 57.5' N  
 Longitude: 144° 43.2' W  
 Water Depth: 234 meters

Lithology: Very soft, soupy, gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Planktic Foraminifers  
 Ostracodes

Ostracode Species:

Adults Juv. %

<u>Bythocythere</u> sp. B	5		41.67
<u>Cytheropteron</u> sp. Q	2	3	41.67
<u>Palmanella limicola</u> (Norman, 1865)	2		16.67

Total Ostracodes 12

EGAL-75-KC

Van Veen-125

Latitude: 59° 59.8' N  
 Longitude: 144° 44.0' W  
 Water Depth: 232 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Planktic Foraminifers  
 Worm Tubes  
 Ostracodes  
 Diatoms

Ostracode Species:

Adults Juv. %

<u>Loxoconcha</u> sp. B	2		33.33
<u>"Leguminocythereis"</u> sp. A	1		16.67
<u>Loxoconcha</u> sp. A		1	16.67
<u>Cytheropteron</u> sp. D	1		16.67
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975		1	16.67

Total Ostracodes 6

EGAL-75-KC            Box Core-127

Latitude:    60° 02.8' N  
Longitude:   144° 43.5' W  
Water Depth: 210 meters

Lithology:      Gray mud.

Organisms:        Calcareous Benthic Foraminifers  
                    Gastropods  
                    Ostracodes  
                    Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Bythocythere</u> sp. B	9	4	35.14
* <u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)	2	6	21.62
<u>Cytheropteron</u> sp. A	7		18.92
<u>Loxoconcha</u> sp. B	3		8.11
<u>Cytheropteron</u> sp. Q		2	5.41
* <u>Krithe</u> sp. A	2		5.41
<u>Cytheropteron</u> sp. C	2		5.41
Total Ostracodes    37			

EGAL-75-KC            Box Core-128

Latitude:    60° 00.6' N  
Longitude:   144° 40.0' W  
Water Depth: 227 meters

Lithology:      Stiff gray mud.

Organisms:        Calcareous Benthic Foraminifers  
                    Agglutinated Benthic Foraminifers  
                    Planktic Foraminifers  
                    Cheiostome Bryozoans  
                    Cyclostome Bryozoans  
                    Ostracodes  
                    Echinoderm Fragments  
                    Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974	14		17.72
<u>Cythere</u> sp. A	8		10.13
<u>Cytheropteron</u> sp. B	2	5	8.86
<u>Cytheropteron</u> sp. A	3	3	7.60
<u>Bythocythere</u> sp. B	5	1	7.60
<u>Loxoconcha</u> sp. A	1	5	7.60
<u>Aurila</u> sp. A		4	5.06
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952	1	3	5.06
<u>Loxoconcha</u> sp. D	1	2	3.80
<u>Cythweromorpha</u> sp. B	2	1	3.80
<u>Paradoxostoma</u> sp. G		3	3.80
<u>Coquimba</u> sp. A		2	2.53
<u>Hemicytherura</u> sp. B	1	1	2.53
<u>Hemicytherura</u> sp. A	2		2.53
<u>Krithe</u> sp. A	1		1.27
<u>Pontocythere</u> sp. A	1		1.27
<u>Cytherura</u> sp. G	1		1.27
<u>Paradoxostoma</u> sp. D		1	1.27
" <u>Australicythere</u> " sp. A		1	1.27
<u>Cluthia</u> sp. A	1		1.27
" <u>Leguminocythereis</u> " sp. A		1	1.27
<u>Cytheropteron</u> sp. X	1		1.27
" <u>Leguminocythereis</u> " sp. D		1	1.27

Total Ostracodes 79

EGAL-75-KC

Shipek-129

Latitude: 60° 04.9' N  
 Longitude: 144° 40.4' W  
 Water Depth: 146 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Proteinaceous Worm Tubes  
 Pelecypods  
 Ostracodes  
 Fish Vertebrae  
 Plant Material

Ostracode Species:	Adults	Juv.	%
" <u>Leguminocythereis</u> " sp. A	9	85	47.96
* <u>Pectocythere</u> sp. D	73	6	40.31
<u>Loxoconcha</u> sp. A	6	6	7.14
<u>Cytheromorpha</u> sp. A	6		3.06
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	2		1.02
<u>Aurila</u> sp. A		1	0.51

Total Ostracodes 196

EGAL-75-KC Shipek-130

Latitude: 60° 07.8' N  
 Longitude: 144° 39.5' W  
 Water Depth: 31 meters

Lithology: Fine sand with muddy matrix.

Organisms: Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron</u> sp. Z	1	2	33.33
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975	2		22.22
<u>Loxoconcha</u> sp. B	2		22.22
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)		1	11.11
<u>Cytherella</u> sp. A		1	11.11

Total Ostracodes 9

EGAL-75-KC            Shipek-132

Latitude:    60° 07.1' N  
Longitude:   144° 31.2' W  
Water Depth: 20 meters

Lithology:      Gray sand.

Organisms:       Ostracodes

Ostracode Species:	Adult	Juv.	%
" <u>Leguminocythereis</u> " sp. B	2	2	44.44
<u>Pectocythere</u> sp. D	1		11.11
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952		1	11.11
<u>Loxoconcha</u> sp. A	1		11.11
<u>Candonia</u> sp.		1	11.11
<u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974		1	11.11
		1	11.11

Total Ostracodes 9

EGAL-75-KC            Shipek-133

Latitude:    60° 03.8' N  
Longitude:   144° 26.2'W  
Water Depth: 17 meters

Lithology:      Fine-grained, gray sand.

Organisms:       Ostracodes

Ostracode Species:	Adult	Juv.	%
<u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974	1		100

Total Ostracodes 1

EGAL-75-KC

Shipek-134

Latitude: 59° 59.0' N  
Longitude: 144° 24.0' W  
Water Depth: 20 meters

Lithology: Fine-grained sand with very slight amount of mud.

Organisms: Ostracodes

Ostracode Species:	Adults	Juv.	%
* <u>Pontocythere</u> sp. A	27	1	22.40
* <u>"Leguminocythereis"</u> sp. B	14	13	21.60
* <u>Pectocythere</u> sp. D	26		20.80
<u>Aurila</u> sp. A	2	9	8.80
<u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974	2	7	7.20
* <u>Loxoconcha</u> sp. A	5	1	4.80
* <u>"Leguminocythereis"</u> sp. A	6		4.80
<u>"Radimella"</u> <u>jollaensis</u> (LeRoy, 1943)	1	2	2.40
<u>Cytheromorpha</u> sp. A	3		2.40
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952	1	1	1.60
<u>Loxoconcha</u> sp. D		1	0.80
<u>Finmarchinella</u> ( <u>Barentsovia</u> ) <u>angulata</u> (Sars, 1866)	1		0.80
<u>Robertsonite</u> <u>tuberculata</u> (Sars, 1865)		1	0.80
<u>Cythere</u> sp. A		1	0.80
Total Ostracodes	125		

EGAL-75-KC

Shipek-138

Latitude: 59° 38.2' N  
Longitude: 145° 50.4' W  
Water Depth: 168 meters

Lithology: Gray mud.

Organisms:

- Calcareous Benthic Foraminifers
- Agglutinated Benthic Foraminifers
- Sponge Spicules
- Pelecypods
- Pteropod
- Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Palmanella limicola</u> (Norman, 1865)		2	22.22
<u>Munseyella</u> sp. A	2		22.22
<u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)	1	1	22.22
<u>Argilloecia</u> sp. A		1	11.11
<u>Eucytherura</u> sp. A	1		11.11
<u>Cytheropteron</u> sp. L		1	11.11
Total Ostracodes	9		

EGAL-75-KC              Van Veen-141

Latitude:    60° 06.8' N  
 Longitude:   146° 14.5' W  
 Water Depth: 71 meters

Lithology:      Gray mud with muddy sand; coarse-grained, poorly-sorted, rounded sand layer of variable thickness; pebbles in mud.

Organisms:

- Calcareous Benthic Foraminifers
- Planktic Foraminifers
- Sponge Spicules
- Bryozoans
- Mollusk Fragments
- Ostracodes
- Echinoderm Fragments

Ostracode Species:	Adults	Juv.	%
<u>"Australicythere"</u> sp. A	26	96	38.49
<u>Pectocythere</u> aff. <u>P. parkerae</u> Swain and Gilby, 1974	32	8	12.62
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	22	11	10.41
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	15	7	6.94
<u>Cytheropteron</u> sp. G	10	4	4.42
<u>Finmarchinella</u> ( <u>Barentosvia</u> ) sp. A	8	3	3.47

Ostracode Species:	Adult	Juv.	%
<u>Ambostracon</u> sp. A	5	5	3.16
" <u>Leguminocythereis</u> " sp. D	4	4	2.52
<u>Cytheropteron</u> sp. H	4	4	2.52
<u>Hemicytherura</u> sp. B	6		1.89
<u>Cytheropteron</u> sp. E	4	2	1.89
<u>Finnmarchinella</u> ( <u>Finnmarchinella</u> )	1	4	1.58
<u>finmarchica</u> (Sars, 1866)			
<u>Acuminocythere</u> sp. A	3	2	1.58
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)		4	1.26
<u>Cytheromorpha</u> sp. B	4		1.26
<u>Eucytherura</u> sp. A	3		0.95
* <u>Argilloecia</u> sp. B	2		0.63
<u>Cytheropteron</u> sp. I	1	1	0.63
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1965)	1	1	0.63
<u>Cytheromorpha</u> sp. A	2		0.63
<u>Coquimba</u> sp. A	1		0.32
<u>Hemicytherura</u> sp. A	1		0.32
<u>Cytheropteron</u> sp. N	1		0.32
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	1		0.32
<u>Pectocythere</u> sp. F	1		0.32
<u>Cytheropteron</u> sp. X	1		0.32
<u>Cytheropteron</u> sp. F	1		0.32
<u>Cytheropteron</u> sp.	1		0.32

Total Ostracodes 317

EGAL-75-KC                  Van Veen-144-Upper

Latitude:        59° 57.3' N  
 Longitude:      146° 19.6' W  
 Water Depth:    64 meters

Lithology:       Sandy mud with shells and gravel.

Organisms:        Calcareous Benthic Foraminifers  
                     Planktic Foraminifers  
                     Sponge Spicules  
                     Agglutinated Worm Tubes  
                     Cheiostome Bryozoans  
                     Brachiopods  
                     Pelecypods  
                     Cirriped Plates  
                     Ostracodes  
                     Echinoderm Fragments  
                     Diatoms

Ostracode Species:	Adults	Juv.	%
* "Australicythere" sp. A	66	284	41.32
* <u>Pectocythere</u> aff. <u>P. parkerae</u> Swain and Gilby, 1974	93	9	12.04
<u>Ambostacon</u> sp. A	50	28	9.21
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	59	16	8.86
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	26	13	4.61
<u>Finmarchinella</u> ( <u>Barentsovia</u> ) sp. A	32	6	4.49
<u>Patagonacythere</u> sp. A	13	9	2.60
<u>Cytheropteron</u> sp. G	14	8	2.60
" <u>Leguminocythereis</u> " sp. D	8	6	1.65
<u>Finmarchinella</u> ( <u>Finmarchinella</u> ) <u>finmarchica</u> (Sars, 1866)	6	8	1.65
<u>Aurila</u> sp. A	3	9	1.42
<u>Hemicytherura</u> sp. B	10	1	1.30
<u>Cytheropteron</u> sp. N	7	4	1.30
<u>Cytheropteron</u> sp. E	3	6	1.06
<u>Acuminocythere</u> sp. A	3	4	0.83
<u>Cytheromorpha</u> sp. B	6	1	0.83
<u>Cytherura</u> sp. F	5		0.59
<u>Pectocythere</u> sp. E	5		0.59
<u>Cytheropteron</u> sp. H	3	1	0.47
<u>Cytheromorpha</u> sp. A	4		0.47
<u>Pectocythere</u> sp. F	3		0.35
<u>Cytheropteron</u> sp. F	3		0.35
<u>Cytheropteron</u> sp. I	2		0.24
<u>Sclerochilus</u> sp. F	1	1	0.24
<u>Loxoconcha</u> sp. E	2		0.24
<u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974	1		0.12
" <u>Acanthocythereis</u> " <u>dunelnensis</u> (Norman, 1865)		1	0.12
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952	1		0.12
<u>Argilloecia</u> sp. B	1		0.12
<u>Sclerochilus</u> sp. G	1		0.12
<u>Sclerochilus</u> sp. C		1	0.12

Total Ostracodes 847

EGAL-75-KC Shipek-145

Latitude: 59° 37.4' N  
Longitude: 146° 09.0' W  
Water Depth: 101 meters

Lithology: Thin layer of soupy, olive-gray mud overlying gray mud.

Organisms:      Calcareous Benthic Foraminifers  
                   Agglutinated Benthic Foraminifers  
                   Proteinaceous Worm Tubes  
                   Pelecypods  
                   Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	4		23.53
<u>Palmanella limicola</u> (Norman, 1865)	4		23.53
<u>Robertsonites tuberculata</u> (Sars, 1865)	3		17.65
<u>Cytheropteron</u> sp. L	3		17.65
<u>Munseyella</u> sp. B	1		5.88
Total Ostracodes 17			

EGAL-75-KC      Shipek-146

Latitude:      59° 35.6' N  
                   Longitude: 145° 54.8' W  
                   Water Depth: 143 meters

Lithology:      Thin layer of soupy, olive-green mud overlying gray mud.

Organisms:      Calcareous Benthic Foraminifers  
                   Agglutinated Benthic Foraminifers  
                   Proteinaceous Worm Tubes  
                   Pteropods  
                   Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Palmanella limicola</u> (Norman, 1865)	2	7	47.37
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)		5	26.32
<u>Loxoconcha</u> sp. B	3		15.79
<u>Cytheropteron</u> sp. Q	1	1	10.53
Total Ostracodes 19			

EGAL-75-KC

Shipek-147

Latitude: 59° 34.2' N  
Longitude: 145° 45.7' W  
Water Depth: 165 meters

Lithology: Soupy, olive-gray mud overlying gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Planktic Foraminifers  
Radiolarians  
Proteinaceous Worm Tubes  
Pelecypods  
Pteropod  
Ostracodes  
Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron</u> sp. L	2	1	30.00
<u>Loxoconcha</u> sp. B	2	1	30.00
<u>Palmanella limicola</u> (Norman, 1865)		1	10.00
<u>Cytheropteron</u> sp. Q		1	10.00
<u>Munseyella</u> sp. B	1		10.00
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	1		10.00

Total Ostracodes 10

EGAL-75-KC

Shipek-149

Latitude: 60° 03.2' N  
Longitude: 145° 34.5' W  
Water Depth: 104 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Planktic Foraminifers  
Proteinaceous Worm Tubes  
Pelecypods  
Ostracodes  
Diatoms

Ostracode Species:	Adult	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	11	30	53.25
<u>Palmanella limicola</u> (Norman, 1865)	5	4	11.69
<u>Eucytherura</u> sp. A	8		10.40
<u>Loxoconcha</u> sp. A	3	2	6.49
<u>Munseyella</u> sp. A	4		5.20
<u>Cytheropteron</u> sp. L	2	1	3.90
<u>Robertsonites tuberculata</u> (Sars, 1865)		3	3.90
<u>Loxoconcha</u> sp. B	2		2.60
<u>Munseyella</u> sp. B	1		1.30
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)		1	1.30

Total Ostracodes 77

EGAL-75-KC Van Veen-150

Latitude: 60° 10.4' N  
 Longitude: 145° 34.5' W  
 Water Depth: 104 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Agglutinated Benthic Foraminifers  
 Planktic Foraminifers  
 Radiolarians  
 Pelecypods  
 Ostracodes  
 Plant Material  
 Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Robertsonites tuberculata</u> (Sars, 1865)	5	32	25.00
<u>Loxoconcha</u> sp. B	22	8	20.27
<u>Cytheropteron</u> sp. A	8	19	18.24
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	2	20	14.87
<u>Cytheropteron</u> sp. Q	5	4	6.08
<u>Bythocythere</u> sp. B	3	1	2.70
<u>Cluthia</u> sp. A	3	1	2.70
<u>Pseudocythere</u> sp. A	2		1.35
<u>Munseyella</u> sp. B	2		1.35
<u>Palmanella limicola</u> (Norman, 1865)	1	1	1.35

Ostracode Species:	Adult	Juv.	%
<u>Cytheropteron</u> sp. K	2		1.35
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	1	1	1.35
<u>Loxoconcha</u> sp. A		1	0.68
<u>Cytherois</u> sp. A	1		0.68
<u>Pectocythere</u> aff. <u>P. parkerae</u> Swain and Gilby, 1974		1	0.68
<u>Cytherura</u> sp. I		1	0.68
<u>Cyclocypris</u> sp.	1		0.68

Total Ostracodes 148

EGAL-75-KC Van Veen-153

Latitude: 60° 12.5' N  
 Longitude: 146° 27.0' W  
 Water Depth: 137 meters

Lithology: Olive-gray, soupy mud overlying homogeneous gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Pelecypods  
 Pteropod  
 Ostracodes  
 Echinoderm Fragments  
 Plant Material  
 Diatoms

Ostracode Species:	Adults	Juv.	%
"Acanthocythereis" <u>dunelmensis</u> (Norman, 1865)	14	135	56.87
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)	8	55	24.05
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	2	11	4.96
<u>Cytheropteron</u> sp. A	5	5	3.82
<u>Bythocythere</u> sp. B	5		1.91
<u>Loxoconcha</u> sp. B	4	1	1.91
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)		4	1.53
" <u>Australicythere</u> " sp. A		3	1.15
<u>Cytherura</u> sp. I		2	0.76
<u>Cluthia</u> sp. A	2		0.76
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957		1	0.38
<u>Cytheromorpha</u> sp. B	1		0.38

Ostracode Species:	Adult	Juv.	%
<u>Hemicytherura</u> sp. A		1	0.38
<u>Xestoleberis</u> sp. B		1	0.38
<u>Cluthia</u> <u>cluthae</u> (Brady, Crosskey and Robertson, 1874)	1		0.38
<u>Finmarchinella</u> ( <u>Finmarchinella</u> ) <u>finmarchica</u> (Sars, 1866)		1	0.38
Total Ostracodes 262			

EGAL-75-KC Van Veen-154

Latitude: 59° 51.4' N  
Longitude: 145° 28.5' W  
Water Depth: 95 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Proteinaceous Worm Tubes  
Pelecypods  
Pteropod  
Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Loxoconcha</u> sp. A	5	5	25.64
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	5	2	17.95
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	1	6	17.95
<u>Munseyella</u> sp. A	5		12.82
<u>Eucytherura</u> sp. A		2	5.13
<u>Cytherura</u> sp. I		2	5.13
<u>Cytheropteron</u> sp. L		2	5.13
<u>Cytheropteron</u> sp. D	2		5.13
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)		1	2.56
<u>Munseyella</u> sp. B	1		2.56
Total Ostracodes 39			

EGAL-75-KC      Van Veen-155

Latitude:    59° 55.2' N  
Longitude:   145° 42.0' W  
Water Depth: 82 meters

Lithology:     Gray mud with pebbles, cobbles, and shells.

Organisms:     Calcareous Benthic Foraminifers  
                  Sponge Spicules  
                  Bryozoans  
                  Proteinaceous Worm Tubes  
                  Pelecypods  
                  Gastropods  
                  Pteropod  
                  Ostracodes  
                  Echinoderms

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron aff. C. latissimum</u> of Neale and Howe (1975)	1	1	33.33
"Australicythere" sp. A		2	33.33
<u>Cytheromorpha</u> sp. B	2		33.33
Total Ostracodes 6			

EGAL-75-KC      Van Veen-157

Latitude:    60° 01.4' N  
Longitude:   146° 08.5' W  
Water Depth: 73 meters

Lithology:     Gray mud with pebbles and cobbles.

Organisms:     Calcareous Benthic Foraminifers  
                  Agglutinated Benthic Foraminifers  
                  Planktic Foraminifers  
                  Cheiostome Bryozoans  
                  Cyclostome Bryozoans  
                  Proteinaceous Worm Tubes  
                  Pelecypods  
                  Gastropods  
                  Scaphopod Fragments  
                  Ostracodes  
                  Echinoderm Fragments

Ostracode Species:	Adults	Juv.	%
<u>Aurila</u> sp. A	20	110	20.12
<u>Ambostacon</u> sp. A	5	33	5.88
" <u>Acanthocythereis</u> " <u>dunelmensis</u>		38	5.88
<u>Pectocythere</u> aff. <u>P. parkerae</u>	20	14	5.26
Swain and Gilby, 1974			
<u>Loxoconcha</u> sp. A	13	16	4.49
<u>Cytheropteron</u> aff. <u>C. latissimum</u>	10	17	4.18
of Neale and Howe (1975)			
<u>Acuminocythere</u> sp. A	3	24	4.18
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)	2	21	3.56
<u>Cytheropteron</u> sp. E	14	8	3.41
<u>Cytheropteron</u> sp. A	11	9	3.10
<u>Cytheromorpha</u> sp. B	10	8	2.79
<u>Cytheropteron</u> sp. N	12	4	2.48
<u>Cytheromorpha</u> sp. A	6	7	2.01
<u>Sclerochilus</u> sp. C	4	9	2.01
<u>Loxoconcha</u> sp. D	6	5	1.70
<u>Pseudocythere</u> sp. A	10	1	1.70
<u>Argilloecia</u> sp. A	6	4	1.55
<u>Pectocythere</u> aff. <u>P. quadrangulata</u>	2	8	1.55
Hanai, 1957			
<u>Hemicytherura</u> sp. B	5	5	1.55
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	4	6	1.55
<u>Paradoxostoma</u> sp. H	7	3	1.55
<u>Xestoleberis</u> sp. B	4	5	1.39
" <u>Australicythere</u> " sp. A		9	1.39
<u>Cytheropteron</u> sp. R	8	1	1.39
<u>Paradoxostoma</u> sp. I	5	4	1.39
<u>Paradoxostoma</u> sp. D	8		1.24
" <u>Radimella</u> " <u>jollaensis</u> (LeRoy, 1943)	1	7	1.24
<u>Cytheropteron</u> sp. F	5	1	0.93
<u>Finmarchinella</u> ( <u>Barentsovia</u> ) sp. A	1	5	0.93
<u>Cluthia</u> <u>cluthae</u> (Brady, Crosskey, and Robertson, 1874)	5	1	0.93
<u>Coquimba</u> sp. A	4	2	0.93
<u>Bythocytheromorpha</u> sp. A	4	1	0.77
* <u>Bairdia</u> sp. A	2	2	0.62
<u>Munseyella</u> sp. A	4		0.62
<u>Cytheropteron</u> sp. I	3	1	0.62
<u>Cytheropteron</u> sp. G	2	1	0.46
<u>Buntonia</u> sp. A		3	0.46
<u>Cytheromorpha</u> sp. E	3		0.46
<u>Sclerochilus</u> sp. E	3		0.46
<u>Argilloecia</u> sp. B		2	0.31
<u>Bythocytheromorpha</u> sp. B	2		0.31
<u>Semicytherura</u> aff. <u>S. undata</u> (Sars, 1865)	2		0.31
<u>Hemicytherura</u> sp. A	2		0.31
<u>Hemicythere</u> sp.	1	1	0.31
<u>Cytheropteron</u> sp. Y		2	0.31
<u>Paracypris</u> sp.	1		0.16

Ostracode Species:	Adult	Juv.	%
<u>Pectocythere</u> sp. F	1		0.16
<u>Bythocythere</u> sp. B		1	0.31
<u>Cytheropteron</u> sp. H	1		0.16
<u>Cytherura</u> sp. H	1		0.16
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952		1	0.16
<u>Elofsonella</u> sp. A	1		0.16
<u>Schizocythere</u> sp. A	1		0.16
<u>Bythocythere</u> sp. A	1		0.16

Total Ostracodes 646

EGAL-75-KC Van Veen-158

Latitude: 60° 06.0' N  
 Longitude: 146° 40.5' W  
 Water Depth: 117 meters

Lithology: Gray mud.  
 Organisms: Calcareous Benthic Foraminifers  
 Planktic Foraminifers  
 Radiolarians  
 Mollusk Fragments  
 Ostracodes  
 Echinoderm Fragments  
 Diatoms

Ostracode Species:	Adults	Juv.	%
"Acanthocythereis" <u>dunelmensis</u> <u>(Norman, 1865)</u>	3	8	36.67
<u>Rboertsonites</u> <u>tuberculata</u> (Sars, 1865)		8	26.67
<u>Cytheropteron</u> aff. <u>C. latissimum</u> <u>of Neale and Howe (1975)</u>	1	2	10.00
" <u>Australicythere</u> " sp. A		3	10.00
<u>Cytheropteron</u> sp. N	1		3.33
<u>Loxoconcha</u> sp. A	1		3.33
<u>Cytheropteron</u> sp. A	1		3.33
<u>Munseyella</u> sp. B	1		3.33
<u>Cytheropteron</u> sp. B		1	3.33

Total Ostracodes 30

EGAL-75-KC

Shipek-159

Latitude: 60° 10.2' N  
 Longitude: 146° 52.1' W  
 Water Depth: 165 meters

Lithology: Thin layer of olive-gray, soupy mud overlying gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Planktic Foraminifers  
 Proteinaceous Worm Tubes  
 Bryozoans  
 Pelecypods  
 Cyprid Barnacles  
 Ostracodes  
 Echinoderm Spine  
 Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Aurila</u> sp. A	18	58	16.27
* <u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)	5	43	10.28
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	2	24	5.57
<u>Cytheropteron</u> sp. A	13	10	4.93
<u>Loxoxoncha</u> sp. A	5	16	4.50
<u>Ambostracon</u> sp. A	1	19	4.28
<u>Cytheropteron</u> sp. N	11	9	4.28
<u>Paradoxostoma</u> sp. G	9	9	3.85
<u>Cytheromorpha</u> sp. B	10	6	3.43
* <u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)	1	15	3.43
" <u>Australicythere</u> " sp. A		13	2.78
<u>Loxoconcha</u> sp. D	7	5	2.57
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	10	1	2.36
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	4	7	2.36
<u>Cytheropteron</u> sp. E	4	6	2.14
<u>Finmarchinella</u> ( <u>Barentsovia</u> ) sp. A	1	8	1.93
* <u>Acuminocythere</u> sp. A	2	7	1.93
<u>Sclerochilus</u> sp. C	4	4	1.71
<u>Xestoleberis</u> sp. B	6	1	1.50
<u>Cytheropteron</u> sp. F	1	6	1.50
<u>Paradoxostoma</u> aff. <u>P. japonicum</u> Schornikov, 1975	3	3	1.29
<u>Cytherois</u> sp. A	5	1	1.29
<u>Hemicytherura</u> sp. A	6		1.29
<u>Cytheromorpha</u> sp. E	5		1.07
<u>Pseudocythere</u> sp. A	4	1	1.07
* <u>Munseyella</u> sp. A	4		0.86
<u>Coquimba</u> sp. A	4	0.86	

Ostracode Species:	Adult	Juv.	%
<u>Paradoxostoma aff. P. brunneatum</u>	1	2	0.64
Schornikov, 1975			
* <u>Bairdia</u> sp.	2	1	0.64
<u>Argilloecia</u> sp. A	1	2	0.64
<u>Cytheropteron</u> sp. G	2	1	0.64
<u>Hemicytherura</u> sp. B	3		0.64
<u>Loxoconcha</u> sp. E	2	1	0.64
<u>Cytheropteron</u> sp. I		3	0.64
<u>Cytheropteron</u> sp. H		3	0.64
<u>Semicytherura</u> aff. <u>S. undata</u>	2		0.43
(Sars, 1865)			
<u>Hemicytherura</u> sp. C	2		0.43
<u>Elofsonia</u> sp. A	2		0.43
<u>Xiphichilus</u> sp.	2		0.43
<u>Pectocythere</u> sp. E	2		0.43
<u>Buntonia</u> sp. A	2		0.43
<u>Bythocytheromorpha</u> sp. B	1		0.21
<u>Xestoleberis</u> sp. A		1	0.21
" <u>Leguminocythereis</u> : sp. D		1	0.21
<u>Cytheropteron</u> sp. Y	1		0.21
<u>Paradoxostoma</u> sp. J	1		0.21
<u>Bythocythere</u> sp. A	1		0.21
<u>Normanicythere</u> sp.		1	0.21
<u>Argilloecia</u> sp. B	1		0.21
<u>Eucytherura</u> sp. A		1	0.21
<u>Semicytherura</u> sp. F		1	0.21
<u>Hemicythere</u> aff. <u>H. quadrinodosa</u>		1	0.21
Schornikov, 1974			

Total Ostracodes 467

EGAL-75-KC

Van Veen-160A

Latitude: 60° 15.0' N  
 Longitude: 146° 31.8' W  
 Water Depth: 37 meters

Lithology: Dark, fine-grained sand.

Organisms: Calcareous Benthic Foraminifers  
 Pelecypods  
 Echinoderm Spines

EGAL-75-KC            Box Core-160C

Latitude:        60° 14.9' N  
Longitude:      146° 32.2' W  
Water Depth:    37 meters

Lithology:        Dark, fine-grained sand.

Organisms:        Calcareous Benthic Foraminifers  
                    Agglutinated and Proteinaceous Worm Tubes  
                    Pelecypods

EGAL-75-KC            Van Veen-161

Latitude:        60° 17.4' N  
Longitude:      146° 23.7' W  
Water Depth:    22 meters

Lithology:        Dark, fine-grained, well-sorted sand.

Organisms:        Calcareous Benthic Foraminifers  
                    Pelecypods  
                    Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Hemicythere aff. H. quadrinodosa</u> Schornikov, 1974	1		50.00
<u>Cythere aff. C. alveolivalva</u> Smith, 1952	1		50.00
Total Ostracodes 2			

EGAL-75-KC            Van Veen-162

Latitude:        60° 19.2' N  
Longitude:      146° 13.2' W  
Water Depth:    24 meters

Lithology:        Dark, fine-grained sand with overlying thin layer of soupy,  
                    gray mud.

Organisms:        Calcareous Benthic Foraminifers  
                    Pelecypods  
                    Ostracodes  
                    Echinoderm Fragments  
                    Plant Material

Ostracode Species:	Adults	Juv.	%
<u>Cytheromorpha</u> sp. B	3	6	33.33
<u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974	3	2	18.52
* <u>Elofsonia</u> sp. A	4		14.82
<u>Pectocythere</u> sp. D	2	1	11.11
<u>Cytherura</u> sp. D	1	2	11.11
<u>Cytheromorpha</u> sp. E	2		7.41
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952		1	3.70
Total Ostracodes 27			

EGAL-75-KC              Van Veen-163

Latitude:    60° 19.5' N  
 Longitude:   146° 07.0' W  
 Water Depth: 22 meters

Lithology:      Dark, fine-grained sand.

Organisms:        Calcareous Benthic Foraminifers  
                     Pelecypods  
                     Ostracodes  
                     Plant Material

Ostracode Species:	Adults	Juv.	%
* <u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974	2		66.67
<u>Elofsonia</u> sp. A	1		33.33
Total Ostracodes 3			

EGAL-75-KC              Van Veen-164A

Latitude:    60° 19.5' N  
 Longitude:   146° 00.0' W  
 Water Depth: 22 meters

Lithology:      Dark, fine-grained sand.

Organisms:        Calcareous Benthic Foraminifers  
                     Plant Debris

EGAL-75-KC

Van Veen-164B

Latitude: 60° 19.5' N  
 Longitude: 146° 00.0' W  
 Water Depth: 22 meters

Lithology: Dark, fine-grained sand.

Organisms: Occasional Calcareous Benthic Foraminifers  
 Pelecypod Fragments  
 Ostracodes  
 Occasional Echinoderm Fragments  
 Plant Fragments

Ostracode Species:

Adults Juv. %

<u>Cytheromorpha</u> sp. B	5	1	46.15
<u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974	2	2	30.77
<u>Elofsonia</u> sp. A	3		23.08

Total Ostracodes 13

EGAL-75-KC

Shipek-165

Latitude: 60° 18.3' N  
 Longitude: 145° 53.5' W  
 Water Depth: 33 meters

Lithology: Dark, fine-grained sand with some gray mud.

Organisms: Rare Calcareous Benthic Foraminifers  
 Pelecypods  
 Ostracodes

Ostracode Species:

Adults Juv. %

* <u>Cytheromorpha</u> sp. B	2	3	41.67
<u>Pectocythere</u> sp. D	2		16.67
* <u>Loxoconcha</u> sp. A	2		16.67
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957		1	8.33
<u>Eucythere</u> sp. A	1		8.33
" <u>Leguminocythereis</u> " sp. A		1	8.33

Total Ostracodes 12

EGAL-75-KC            Shipek-166

Latitude:        60° 17.7' N  
Longitude:      145° 45.6' W  
Water Depth:    20 meters

Lithology:       Dark, fine-grained sand.

Organisms:       Calcareous Benthic Foraminifers  
                    Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Candonia</u> sp.		1	25.00
<u>Elofsonia</u> sp. A	1		25.00
<u>Aurilla</u> sp. A		1	25.00
<u>Cytheromorpha</u> sp. B	1		25.00
Total Ostracodes	4		

EGAL-75-KC            Shipek-167

Latitude:        60° 16.4' N  
Longitude:      145° 38.4' W  
Water Depth:    26 meters

Lithology:       Dark, fine-grained sand.

Organisms:       Calcareous Benthic Foraminifers  
                    Mollusk Fragments  
                    Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>"Leguminocythereis"</u> sp. A		1	100
Total Ostracodes	1		

EGAL-75-KC            Shipek-169

Latitude:        60° 17.7' N  
Longitude:      145° 50.7' W  
Water Depth:    35 meters

Lithology: Dark, fine-grained sand with some clay galls at surface.

Organisms: Calcareous Benthic Foraminifers  
Pelecypods  
Echinoderm Fragments

EGAL-75-KC Box Core-170B

Latitude: 60° 16.9' N  
Longitude: 145° 42.0' W  
Water Depth: 20 meters

Lithology: Dark, fine-grained, well-sorted sand overlying gray mud.

Organisms: Calcareous Benthic Foraminifers  
Pelecypods  
Ostracodes  
Echinoderm Fragments  
Plant Fragments

Ostracode Species:	Adults	Juv.	%
<u>Cytheromorpha</u> sp. B	2		50.00
<u>Pectocythere</u> sp. D	1		25.00
<u>Loxoconcha</u> sp. A		1	25.00
Total Ostracodes 4			

EGAL-75-KC Shipek-172

Latitude: 60° 13.5' N  
Longitude: 145° 21.2' W  
Water Depth: 18 meters

Lithology: Dark, fine-grained sand.

Organisms: Barren of any faunal or floral remains.

EGAL-75-KC Shipek-174

Latitude: 60° 09.6' N  
Longitude: 145° 06.4' W  
Water Depth: 35 meters

Lithology: Gray mud with fine-grained sand.

Organisms: Ostracodes

Ostracode Species:	Adults	Juv.	%
" <u>Leguminocythereis</u> " sp. A	3	3	37.50
<u>Pectocythere</u> sp. D	5		31.25
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957	2		12.50
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)		1	6.25
<u>Loxoconcha</u> sp. A		1	6.25
<u>Cytheropteron</u> sp. R	1		6.25
Total Ostracodes	16		

EGAL-75-KC Shipek-175

Latitude: 60° 09.4' N

Longitude: 145° 00.0' W

Water Depth: 33 meters

Lithology: Gray mud with very thin, surficial layer of sand.

Organisms: Calcareous Benthic Foraminifers  
Pelecypods  
Ostracodes  
Plant Material

Ostracode Species:	Adults	Juv.	%
<u>Cytheromorpha</u> sp. A	1		50.00
<u>Cytheromorpha</u> sp. B		1	50.00
Total Ostracodes	2		

EGAL-75-KC Box Core-179B

Latitude: 60° 14.8' N

Longitude: 145° 26.9' W

Water Depth: 18 meters

Lithology: Fine-grained sand with thin clay layer at surface.

Organisms: Pelecypod Fragments

EGAL-75-KC Shipek-181

Latitude: 60° 01.0' N  
Longitude: 144° 24.0' W  
Water Depth: 33 meters

Lithology: Gray mud with slight amount of sand.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated and Proteinaceous Worm Tubes  
Cheiostome Bryozoans  
Pelecypods  
Gastropods  
Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Pectocythere aff. P. quadrangulata</u> Hanai, 1957	141	55	24.17
* <u>Robertsonites tuberculata</u> (Sars, 1865)	11	150	19.85
<u>Loxoconcha</u> sp. A	48	36	10.36
* <u>Buntonia</u> sp. A	73	8	9.99
* <u>Cytheromorpha</u> sp. B	48	7	6.78
<u>Palmanella limicola</u> (Norman, 1865)	22	11	4.07
<u>Hemicythere aff. H. quadrinodosa</u> Schornikov, 1975	5	27	3.95
" <u>Leguminocythereis</u> " sp. A		26	3.21
<u>Cytheromorpha</u> sp. A	24		2.96
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	2	21	2.84
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952	5	10	1.85
<u>Aurila</u> sp. A	1	13	1.73
<u>Cythere</u> sp. A	2	9	1.36
<u>Astrostracon</u> sp. A		11	1.36
* <u>Pectocythere</u> sp. D	5	3	0.99
<u>Loxoconcha</u> sp. D	5	1	0.74
<u>Loxoconcha</u> sp. E	3	2	0.62
<u>Pontocythere</u> sp. A	5		0.62
<u>Cytheropteron</u> sp. I	4		0.49
* <u>Argilloecia</u> sp. A	3		0.37
<u>Hemicytherura</u> sp. A		2	0.25
<u>Elofsonia</u> sp. A	2		0.25
<u>Cytherura</u> sp. F	2		0.25
<u>Argilloecia</u> sp. B	2		0.25
<u>Cytheropteron</u> sp.		1	0.12

Ostracode Species:	Adult	Juv.	%
<u>Cytherois</u> sp. A	1		0.12
<u>paracyprid</u>	1		0.12
<u>Paradoxostoma</u> sp. G	1		0.12
<u>Acuminocythere</u> sp. A	1		0.12
<u>Paradoxostoma</u> sp. B	1		0.12
Total Ostracodes	811		

EGAL-75-KC              Van Veen-203

Latitude:    59° 32.5' N  
 Longitude:   144° 36.4' W  
 Water Depth: Not Determined

Lithology:        Gray, gravelly mud.  
 Organisms:        Calcareous Benthic Foraminifers  
                     Agglutinated Benthic Foraminifers  
                     Sponge Spicules  
                     Cheilostome Bryozoans  
                     Pelecypods  
                     Echinoderm Spines

EGAL-75-KC              Van Veen-219

Latitude:    59° 36.3' N  
 Longitude:   144° 17.4' W  
 Water Depth: 475 meters

Lithology:        Gray mud.  
 Organisms:        Radiolarians  
                     Sponge Spicules  
                     Ostracodes  
                     Echinoderm Fragments

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron</u> sp. Q	1		100
Total Ostracodes	1		

EGAL-75-KC

Shipek-221

Latitude: 59° 50.1' N  
 Longitude: 144° 27.4' W  
 Water Depth: 29 meters

Lithology: Olive-sandy mud overlying gray clay with some shell fragments.

Organisms: Calcareous Benthic Foraminifers  
 Planktic Foraminifers  
 Cheiostome Bryozoans  
 Pelecypods  
 Gastropods  
 Ostracodes  
 Echinoderm Spines  
 Fish Parts  
 Plant Debris

Ostracode Species:	Adults	Juv.	%
* <u>Pectocythere</u> sp. D	224	38	34.79
* <u>Loxoconcha</u> sp. A	95	125	29.22
" <u>Leguminocythereis</u> " sp. A	5	63	9.03
* <u>Cytheromorpha</u> sp. B	40	22	8.23
* <u>Pectocythere</u> aff. <u>P. parkerae</u> Swain and Gilby, 1974	58	2	7.97
<u>Cytheromorpha</u> sp. A	39	2	5.44
* <u>Cytheromorpha</u> sp. E	17		2.26
* <u>Aurila</u> sp. A	2	3	0.66
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975	4	1	0.66
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957		3	0.40
<u>Cytheropteron</u> sp. A	1	2	0.40
<u>Cythere</u> sp. A		2	0.27
" <u>Leguminocythereis</u> " sp. B		2	0.27
<u>Cythere</u> aff. <u>C. alveolivalva</u> Smith, 1952	1		0.13
<u>Palmanella limicola</u> (Norman, 1865)		1	0.13
<u>Finmarchinella</u> ( <u>Barentsovia</u> ) sp. A		1	0.13
<u>Baffinicythere emarginata</u> (Sars, 1865)		1	0.13
<u>Loxoconcha</u> sp. F	1		0.13

Total Ostracodes 753

EGAL-75-KC

Box Core-223

Latitude: 59° 52.4' N  
Longitude: 144° 18.7' W  
Water Depth: 51 meters

Lithology: Muddy, fine-grained sand.

Organisms: Ostracodes

Ostracode Species:	Adults	Juv.	%
* <u>"Leguminocythereis"</u> sp. A	13	104	43.01
* <u>Pectocythere</u> sp. D	103	4	39.34
<u>Loxoconcha</u> sp. A	14	28	15.44
<u>Cytheropteron</u> sp. A	2	1	1.10
" <u>Leguminocythereis</u> " sp. B		2	0.74
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957		1	0.37

Total Ostracodes 272

EGAL-75-KC

Shipek-225

Latitude: 59° 46.2' N  
Longitude: 144° 11.5' W  
Water Depth: 101 meters

Lithology: Gray mud.

Organisms: Pelecypods  
Ostracodes

Ostracode Species:	Adults	Juv.	%
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	3		42.86
<u>Palmanella limicola</u> (Norman, 1865)	2		28.57
<u>Robertsonites tuberculata</u> (Sars, 1865)	1		14.29
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975	1		14.29

Total Ostracodes 7

EGAL-75-KC

Shipek-227

Latitude: 59° 39.4' N  
Longitude: 144° 04.9' W  
Water Depth: 161 meters

Lithology: Very stiff, olive-gray mud with subrounded pebbles and shells.

Organisms: Radiolarians  
Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron aff. C. latissimum</u> of Neale and Howe (1975)	3	2	55.56
<u>Cytheropteron</u> sp. G	2	1	33.33
<u>Cytheropteron</u> sp. L		1	11.11
Total Ostracodes 9			

EGAL-75-KC

Van Veen-229

Latitude: 59° 34.0' N  
Longitude: 144° 01.2' W  
Water Depth: 1189 meters

Lithology: Stiff, gray-green mud.

Organisms: Radiolarians  
Sponge Spicules  
Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865) unidentifiable fragment	1		100
Total Ostracodes 1			

EGAL-75-KC            Van Veen-231

Latitude:    59° 56.9' N  
Longitude:   144° 09.7' W  
Water Depth: 33 meters

Lithology:      Gray sand.

Organisms:        Calcareous Benthic Foraminifers  
                    Bryozoans  
                    Ostracodes  
                    Echinoderm Fragments  
                    Ophiuroids

Ostracode Species:	Adults	Juv.	%
<u>"Leguminocythereis"</u> sp. A	2		100
Total Ostracodes	2		

EGAL-75-KC            Van Veen-232

Latitude:    59° 57.25' N  
Longitude:   144° 09.9' W  
Water Depth: 49 meters

Lithology:      2-3 cm of green-gray sand overlying gray mud, with sand layers.

Organisms:        Pelecypod Fragments  
                    Ostracodes  
                    Echinoderm Fragment  
                    Plant Fragments  
                    Carbonaceous Plant Fragments

Ostracode Species:	Adults	Juv.	%
<u>Pectocythere</u> sp. D	11		68.75
<u>Loxoconcha</u> sp. A	1	1	12.50
<u>Cytheromorpha</u> sp. A	2		12.50
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u>		1	6.25
Neale and Howe, 1975			
Total Ostracodes	16		

EGAL-75-KC

Van Veen-233

Latitude: 59° 51.6' N  
Longitude: 143° 53.25' W  
Water Depth: 106 meters

Lithology: Olive-green mud overlying gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Radiolarians  
Proteinaceous Worm Tubes  
Pelecypods  
Gastropods  
Ostracodes  
Echinoderm Fragments

Ostracode Species:	Adults	Juv.	%
* <u>"Acanthocythereis dunelmensis</u> (Norman, 1865)	3	6	37.50
* <u>Palmanella limicola</u> (Norman, 1865)		9	37.50
* <u>Cytheropteron</u> sp. Q	2		8.33
<u>Cytheropteron</u> sp. A	1	1	8.33
<u>Pectocythere</u> aff. <u>P. quadrangulata</u> Hanai, 1957		1	4.17
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975		1	4.17

Total Ostracodes 24

EGAL-75-KC

Van Veen-237

Latitude: 59° 51.7' N  
Longitude: 143° 42.5' W  
Water Depth: 225 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
Agglutinated Benthic Foraminifers  
Planktic Foraminifers  
Proteinaceous Worm Tubes  
Pelecypods  
Ostracodes  
Echinoderm Spines  
Plant Debris  
Diatoms

Ostracode Species:	Adults	Juv.	%
* <u>Cluthia cluthae</u> (Brady, Crosskey, and Robertson, 1874)	2		66.67
<u>Palmanella limicola</u> (Norman, 1865)	1		33.33
Total Ostracodes 3			

EGAL-75-KC              Van Veen-239

Latitude:    59° 55.6' N  
 Longitude:   143° 32.4' W  
 Water Depth: 252 meters

Lithology:      Gray mud.

Organisms:        Calcareous Benthic Foraminifers  
                     Planktic Foraminifers  
                     Radiolarians  
                     Scaphopod Fragments  
                     Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Palmanella limicola</u> (Normanm, 1865)	1		100
Total Ostracodes 1			

EGAL-75-KC              Van Veen-241

Latitude:    59° 43.2' N  
 Longitude:   143° 26.6' W  
 Water Depth: 311 meters

Lithology:      Olive-gray mud.

Organisms:        Calcareous Benthic Foraminifers  
                     Agglutinated Benthic Foraminifers  
                     Radiolarians  
                     Sponge Spicules  
                     Fish Parts  
                     Diatoms

EGAL-75-KC              Van Veen-242

Latitude:    59° 39.7' N  
Longitude:   143° 22.9' W  
Water Depth: 289 meters

Lithology:      Olive-gray mud.

Organisms:        Calcareous Benthic Foraminifers

EGAL-75-KC              Van Veen-243

Latitude:    59° 34.9' N  
Longitude:   143° 19.9' W  
Water Depth: 238 meters

Lithology:      Olive-gray mud with sand, gravel and cobbles.

Organisms:        Calcareous Benthic Foraminifers  
                    Agglutinated Benthic Foraminifers  
                    Sponge Spicules  
                    Cheilostome Bryozoans  
                    Pelecypods  
                    Scaphopods  
                    Ophiuroid Parts  
                    Echinoderm Fragments and Spines

EGAL-75-KC              Van Veen-249

Latitude:    59° 58.4' N  
Longitude:   143° 23.0' W  
Water Depth: 152 meters

Lithology:      Gray mud.

Organisms:        Pelecypods  
                    Ostracodes

Ostracode Species:	Adults	Juv.	%
"Acanthocythereis" <u>dunelmensis</u> (Norman, 1865)	1	6	41.18
* <u>Palmanella limicola</u> (Norman, 1865)	5		29.41
<u>Buntonia</u> sp. A	2		11.76
<u>Robertsonites tuberculata</u> (Sars, 1865)		2	11.76
<u>Cytheropteron</u> sp. D	1		5.88

Total Ostracodes 17

EGAL-75-KC            Shipek-254A

Latitude:        59° 31.3' N  
 Longitude:      142° 39.5' W  
 Water Depth:    403 meters

Lithology:       Muddy gravel.

Organisms:       Calcareous Benthic Foraminifers  
 Agglutinated Benthic Foraminifers  
 Sponge Spicules

EGAL-75-KC        Van Veen-258

Latitude:        59° 57.5' N  
 Longitude:      142° 412.0' W  
 Water Depth:    108 meters

Lithology:       About 1-2 cm of olive-green, soupy mud overlying gray,  
 mottled mud.

Organisms:       Calcareous Benthic Foraminifers  
 Planktic Foraminifers  
 Agglutinated Worm Tubes  
 Pelecypods  
 Ostracodes  
 Echinoderm Fragments  
 Plant Debris

Ostracode Species:	Adults	Juv.	%
<u>Palmanella limicola</u> (Norman, 1865)	6	16	37.93
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)	2	8	17.24
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	2	3	8.62
<u>Munseyella</u> sp. A	5		8.62
<u>Eucytherura</u> sp. A	2	1	5.17
<u>Pectocythere</u> sp. D		2	3.45
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)		2	3.45
<u>Cytheropteron</u> sp. D	1	1	3.45
<u>Cytheromorpha</u> sp. B		1	1.72

Ostracode Species:	Adult	Juv.	%
<u>Pectocythere aff. P. quadrangulata</u> Hanai, 1957		1	1.72
<u>Cytheropteron</u> sp. G	1		1.72
<u>Cytheropteron</u> sp. A		1	1.72
<u>Paradoxostoma</u> sp. I		1	1.72
<u>Loxoconcha</u> sp. B	1		1.72
<u>Buntonia</u> sp. A		1	1.72
Total Ostracodes 58			

EGAL-75-KC              Shipek-264

Latitude: 59° 49.5' N  
 Longitude: 142° 30.0' W  
 Water Depth: 134 meters

Lithology: Gray mud.

Organisms:              Calcareous Benthic Foraminifers  
                           Agglutinated Benthic Foraminifers  
                           Planktic Foraminifers  
                           Pelecypods  
                           Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	3	16	45.24
<u>Palmanella limicola</u> (Norman, 1865)	2	3	11.90
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	5		11.90
<u>Robertsonites</u> <u>tuberculata</u> (Sars, 1865)	1	3	9.52
<u>Cytheropteron</u> sp. L	1	2	7.14
<u>Munseyella</u> sp. A	2		4.76
<u>Cytheropteron</u> sp. G	2		4.76
<u>Loxoconcha</u> sp. A		1	2.38
<u>Paradoxostoma</u> sp. B		1	2.38
Total Ostracodes 42			

EGAL-75-KC            Shipek-265

Latitude:        59° 46.2' N  
Longitude:      142° 29.9' W  
Water Depth:    181 meters

Lithology:       Olive-gray mud with pebbles.

Organisms:       Calcareous Benthic Foraminifers  
                  Agglutinated Benthic Foraminifers  
                  Planktic Foraminifers  
                  Proteinaceous Worm Tubes  
                  Pelecypods  
                  Ostracodes  
                  Echinoderm Fragments

Ostracode Species:	Adults	Juv.	%
<u>Robertsonites tuberculata</u> (Sars, 1865)		2	66.67
<u>Palmanella limicola</u> (Norman, 1865)	1		33.33
Total Ostracodes	3		

EGAL-75-KC            Shipek-266

Latitude:        59° 42.5' N  
Longitude:      142° 34.0' W  
Water Depth:    262 meters

Lithology:       Olive mud with cobbles and gravel.

Organisms:       Calcareous Benthic Foraminifers  
                  Agglutinated Benthic Foraminifers  
                  Planktic Foraminifers  
                  Sponge Spicules  
                  Agglutinated and Proteinaceous Worm Tubes  
                  Ostracodes  
                  Echinoderm Spines and Fragments  
                  Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Eucytherura</u> sp. A		1	100
Total Ostracodes	1		

EGAL-75-KC

Shipek-268

Latitude: 59° 40.7' N  
 Longitude: 142° 21.6' W  
 Water Depth: 174 meters

Lithology: Olive-gray mud layer overlying gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Agglutinated Benthic Foraminifers  
 Planktic Foraminifers  
 Radiolarians  
 Sponge Spicules  
 Proteinaceous Worm Tubes  
 Pelecypods  
 Ostracodes  
 Echinoderm Debris  
 Plant Debris

Ostracode Species:	Adults	Juv.	%
<u>Eucytherura</u> sp. A	7	5	38.71
<u>"Acanthocythereis"</u> <u>dunelmensis</u> (Norman, 1865)		8	25.81
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	1	3	12.90
<u>Krithe</u> sp A		2	6.45
<u>Cytheropteron</u> sp. L	1	1	6.45
<u>Palmanella</u> <u>limicola</u> (Norman, 1865)	1		3.23
<u>Sclerochilus</u> sp. C	1		3.23
<u>Munseyella</u> sp. A	1		3.23
Total Ostracodes	31		

EGAL-75-KC

Van Veen-284

Latitude: 59° 50.0' N  
 Longitude: 142° 14.2' W  
 Water Depth: 86 meters

Lithology: Gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Pelecypods  
 Ostracodes  
 Plant Debris

Ostracode Species:	Adults	Juv.	%
* <u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	25	43	30.09
<u>Robertsonites tuberculata</u> (Sars, 1865)		50	22.12
* <u>Palmanella limicola</u> (Norman, 1865)	20	15	15.49
* <u>Pectocythere aff. P. quadrangulata</u> Hanai, 1957	26	4	13.27
<u>Buntonia</u> sp. A	8	2	4.43
<u>Eucytherura</u> sp. A	7		3.10
<u>Cytheropteron</u> sp. E	2	4	2.66
<u>Cytheropteron</u> aff. <u>C. latissimum</u> of Neale and Howe (1975)	5		2.21
<u>Munseyella</u> sp. A	4		1.77
<u>Cytheropteron</u> sp. G	2		0.89
<u>Cytheropteron</u> sp. H		2	0.89
<u>Cytheromorpha</u> sp. D	2		0.89
<u>Cytheropteron</u> sp. F	1		0.44
<u>Pectocythere</u> aff. <u>P. parkerae</u> Swain and Gilby, 1974	1		0.44
<u>Cytheropteron</u> sp. D	1		0.44
<u>Cytheropteron</u> sp. A	1		0.44

Total Ostracodes 226

EGAL-75-KC Shipek-294

Latitude: 59° 48.7' N  
Longitude: 141° 25.0' W  
Water Depth: 29 meters

Lithology: Fine-grained, gray sand.

Organisms: Calcareous Benthic Foraminifers  
Pelecypods  
Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Pectocythere</u> sp. D	1		100
Total Ostracodes 1			

EGAL-75-KC

Van Veen-306

Latitude: 59° 30.4' N  
 Longitude: 141° 30.0' W  
 Water Depth: 161 meters

Lithology: Gray-green mud.

Organisms: Calcareous Benthic Foraminifers  
 Agglutinated Benthic Foraminifers  
 Planktic Foraminifers  
 Radiolarians  
 Proteinaceous Worm Tubes  
 Pelecypods  
 Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Palmanella limicola</u> (Norman, 1865)	6	3	37.50
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)		6	25.00
<u>Krithe</u> sp. A	3	2	20.83
<u>Munseyella</u> sp. A	4		16.67
Total Ostracodes	24		

EGAL-75-KC

Van Veen-307

Latitude: 59° 28.9' N  
 Longitude: 141° 27.8' W  
 Water Depth: 165 meters

Lithology: Thin layer of green mud overlying gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Ostracodes

Ostracode Species:	Adults	Juv.	%
<u>Palmanella limicola</u> (Norman, 1865)	7	6	38.24
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)		9	26.47
<u>Krithe</u> sp. A		5	14.71
<u>Munseyella</u> sp. A	5		14.71
<u>Cytheropteron</u> sp. Q	1	1	5.88
Total Ostracodes	34		

EGAL-75-KC              Van Veen-310

Latitude:    59° 22.1' N  
Longitude:   141° 09.5' W  
Water Depth: 298 meters

Lithology:      Green, muddy, fine-grained sand.

Organisms:      Calcareous Benthic Foraminifers  
                    Agglutinated Benthic Foraminifers  
                    Planktic Foraminifers  
                    Radiolarians  
                    Sponge Spicules  
                    Worm Tubes  
                    Gastropods  
                    Ophiuroids  
                    Echinoderm Fragments and Spines  
                    Diatoms

EGAL-75-KC              Van Veen-326

Latitude:    59° 24.6' N  
Longitude:   140° 14.5' W  
Water Depth: 183 meters

Lithology:      Gray mud.

Organisms:      Calcareous Benthic Foraminifers  
                    Agglutinated Benthic Foraminifers  
                    Planktic Foraminifers  
                    Radiolarians  
                    Sponge Spicules  
                    Proteinaceous Worm Tubes  
                    Pelecypods  
                    Ostracodes  
                    Echinoderm Fragments  
                    Diatoms

Ostracode Species:	Adults	Juv.	%
<u>"Acanthocythereis" dunelmensis</u> (Norman, 1865)	6		46.15
<u>Palmanella limicola</u> (Norman, 1865)	3		23.08
<u>Krithe</u> sp. A		2	15.39
<u>Cytheropteron</u> sp. L	1		7.69
<u>Cytherois</u> sp. A	1		7.69
Total Ostracodes	13		

EGAL-75-KC

Van Veen-331

Latitude: 59° 56.1' N  
 Longitude: 143° 53.4' W  
 Water Depth: 66 meters

Lithology: Olive-green mud overlying gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Agglutinated Benthic Foraminifers  
 Planktic Foraminifers  
 Pelecypods  
 Ostracodes  
 Plant Debris

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron</u> sp. A	10	13	32.86
<u>Loxoconcha</u> sp. A	5	10	21.43
<u>Palmanella limicola</u> (Normanm, 1865)	5	5	14.29
" <u>Leguminocythereis</u> " sp. A		6	8.57
<u>Pectocythere</u> sp. D	3	1	5.71
" <u>Acanthocythereis</u> " <u>dunelmensis</u> (Norman, 1865)		2	2.86
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975		2	2.86
<u>Hemicytherura</u> sp. B			
<u>Finmarchinella</u> ( <u>Barentsovia</u> ) sp. A	1		1.43
<u>Hemicythere</u> aff. <u>H. quadrinodosa</u> Schornikov, 1974		1	1.43
<u>Paradoxostoma</u> sp. B	1		1.43
<u>Paradoxostoma</u> sp. G	1		1.43
<u>Cytheromorpha</u> sp. D	1		1.43
<u>Cluthia</u> sp. A		1	1.43
Total Ostracodes	70		

EGAL-75-KC

Van Veen-334

Latitude: 59° 37.6' N  
 Longitude: 143° 34.0' W  
 Water Depth: 145 meters

Lithology: Pebbly mud.

Organisms: Sponge Spicules

EGAL-75-KC            Shipek-348

Latitude:        59° 31.8' N  
Longitude:      142° 24.8' W  
Water Depth:    631 meters

Lithology:        Homogeneous olive-gray mud.

Organisms:        Calcareous Benthic Foraminifers  
                     Planktic Foraminifers  
                     Radiolarians  
                     Sponge Spicules  
                     Proteinaceous Worm Tubes  
                     Ophiuroid Plates  
                     Diatoms

EGAL-75-KC            Shipek-349

Latitude:        59° 32.6' N  
Longitude:      142° 30.5' W  
Water Depth:    714 meters

Lithology:        Homogeneous olive-green mud.

Organisms:        Calcareous Benthic Foraminifers  
                     Planktic Foraminifers  
                     Radiolarians  
                     Sponge Spicules  
                     Proteinaceous Worm Tubes  
                     Pelecypods  
                     Ophiuroid Plates  
                     Diatoms

EGAL-75-KC            Van Veen-350

Latitude:        59° 34.3' N  
Longitude:      142° 33.0' W  
Water Depth:    578 meters

Lithology:        Very soft, olive mud.

Organisms:        Calcareous Benthic Foraminifers  
                     Planktic Foraminifers  
                     Sponge Spicules  
                     Proteinaceous Worm Tubes  
                     Pelecypods  
                     Fish Scales  
                     Diatoms

EGAL-75-KC            Van Veen-351

Latitude:    59° 35.0' N  
Longitude:   142° 44.3' W  
Water Depth: 615 meters

Lithology:      Green mud.

Organisms:        Calcareous Benthic Foraminifers  
                    Agglutinated Benthic Foraminifers  
                    Planktic Foraminifers  
                    Radiolarians  
                    Sponge Spicules  
                    Proteinaceous Worm Tubes  
                    Echinoderm Fragments  
                    Ophiuroid Spines  
                    Diatoms

EGAL-75-KC            Shipek-405

Latitude:    59° 52.4' N  
Longitude:   141° 37.5' W  
Water Depth: 24 meters

Lithology:      Gray mud with numerous pebbles.

Organisms:        Rare Benthic Foraminifers  
                    Pelecypod Fragments

EGAL-75-KC            Shipek-406

Latitude:    59° 53.0' N  
Longitude:   141° 36.5' W  
Water Depth: 26 meters

Lithology:      Very stiff gray clay with some pebbles.

Organisms:        Calcareous Benthic Foraminifers  
                    Pelecypods  
                    Ostracodes

---

Ostracode Species:	Adults	Juv.	%
--------------------	--------	------	---

---

<u>Cytheropteron</u> sp. A	1		100
----------------------------	---	--	-----

Total Ostracodes	1
------------------	---

EGAL-75-KC	Shipek-407
Latitude:	59° 52.62' N
Longitude:	141° 35.25' W
Water Depth:	35 meters
Lithology:	Mud with encrusting organisms at surface, with cobbles, gravel and sand.
Organisms:	Calcareous Benthic Foraminifers Bryozoans Pelecypods Cirriped Plates
EGAL-75-KC	Shipek-409
Latitude:	59° 51.7' N
Longitude:	141° 32.2' W
Water Depth:	35 meters
Lithology:	Fine-grained, dark sand.
Organisms:	Pelecypod Fragments
EGAL-75-KC	Shipek-412
Latitude:	59° 53.61' N
Longitude:	141° 35.15' W
Water Depth:	40 meters
Lithology:	Dark gray clay with thin veneer of pebbles and shell fragments.
Organisms:	Calcareous Benthic Foraminifers Cheiostome Bryozoans Pelecypods Cirriped Plates
EGAL-75-KC	Shipek-414
Latitude:	59° 55.99' N
Longitude:	141° 37.6' W
Water Depth:	26 meters
Lithology:	Stiff gray clay overlain by veneer of pebbles.

Organisms:      **Calcareous Benthic Foraminifers**  
                  **Pelecypod Fragments**  
                  **Echinoderm Spine**

EGAL-75-KC      Shipek-415

Latitude:        59° 55.22' N  
Longitude:       141° 36.45' W  
Water Depth:     31 meters

Lithology:       Olive-green mud with pebbles and cobbles overlying gray mud.

Organisms:       **Calcareous Benthic Foraminifers**  
                  **Bryozoans**  
                  **Pelecypods**

EGAL-75-KC      Shipek-422

Latitude:        59° 55.8' N  
Longitude:       141° 35.6' W  
Water Depth:     68 meters

Lithology:       Gray mud with some sand.

Organisms:       **Calcareous Benthic Foraminifers**  
                  **Pelecypods**  
                  **Ostracodes**  
                  **Diatoms**

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron</u> sp. A	23	18	58.57
<u>Cytheropteron</u> sp. N	7		10.00
<u>Hemicytherura</u> sp. A	2	5	10.00
<u>Loxoconcha</u> sp. D	2	2	5.71
<u>Cytheromorpha</u> sp. B	2		2.86
<u>Cytherura</u> sp. F	2		2.86
<u>Paradoxostoma</u> sp. H	2		2.86
<u>Hemicytherura</u> sp. B	1		1.43
<u>Paradoxostoma</u> sp. G	1		1.43
<u>Cytheropteron</u> sp. F	1		1.43
<u>Loxoconcha</u> sp. A		1	1.43
<u>Cytheropteron</u> sp. E		1	1.43
Total Ostracodes	70		

EGAL-75-KC            Shipek-423

Latitude:    59° 55.5' N  
Longitude:   141° 35.9' W  
Water Depth: 27 meters

Lithology:      Stiff gray clay with coarse sand and cobbles at surface.

Organisms:       Rare Pelecypod Fragments

EGAL-75-KC            Shipek-425

Latitude:    59° 56.7' N  
Longitude:   141° 35.1' W  
Water Depth: 59 meters

Lithology:      Green soupy mud overlying gray mud.

Organisms:       Rare Calcareous Benthic Foraminifers  
                  Pelecypods  
                  Gastropod  
                  Ostracodes  
                  Plant Debris

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron</u> sp. A	3	5	72.73
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975	1		9.09
<u>Candonia</u> sp.		1	9.09
<u>Bythocythere</u> sp. B	1		9.09
Total Ostracodes	11		

EGAL-75-KC            Shipek-427

Latitude:    59° 55.45' N  
Longitude:   141° 32.7' W  
Water Depth: 71 meters

Lithology:      Olive-green mud overlying gray-green mud.

Organisms:       Occasional Calcareous Benthic Foraminifers  
                  Pelecypod  
                  Ostracodes  
                  Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron</u> sp. A	1		100
Total Ostracodes 1			

EGAL-75-KC            Shipek-428

Latitude:        59° 54.7' N  
 Longitude:      141° 30.1' W  
 Water Depth:    49 meters

Lithology:       Dark gray, soupy sand with much organic material.

Organisms:       Rare Calcareous Benthic Foraminifers  
 Pelecypods  
 Gastropods  
 Ostracodes  
 Plant Debris  
 Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Loxoconcha</u> sp. A	2		28.57
<u>Cytheromorpha</u> sp. B	1		14.29
<u>Cytheropteron</u> sp. I	1		14.29
<u>Cytherura</u> sp. D	1		14.29
<u>Cythere</u> sp. A	1		14.29
<u>Cytheropteron</u> aff. <u>C. nodoscalatum</u> Neale and Howe, 1975	1		14.29
Total Ostracodes 7			

EGAL-75-KC            Shipek-429

Latitude:        59° 55.5' N  
 Longitude:      141° 30.6' W  
 Water Depth:    60 meters

Lithology:       Green soupy mud overlying gray-green, clayey sand.

Organisms:       Pelecypods  
 Ostracodes  
 Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Hemicytherura</u> sp. C	1		100
Total Ostracodes 1			

EGAL-75-KC              Shipek-431

Latitude:    59° 56.5' N  
 Longitude:   141° 33.3' W  
 Water Depth: 59 meters

Lithology:        Olive-green mud overlying gray mud.

Organisms:        Calcareous Benthic Foraminifers  
                   Pelecypods  
                   Ostracodes  
                   Plant Debris

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron</u> sp. A	4		50.00
<u>Cytheropteron</u> sp. E	2		25.00
<u>Cytheropteron</u> aff. <u>C. nodosoalatum</u> Neale and Howe, 1975	1		12.50
<u>Pseudocythere</u> sp. A	1		12.50
Total Ostracodes 8			

EGAL-75-KC              Shipek-432

Latitude:    59° 57.2' N  
 Longitude:   141° 31.6' W  
 Water Depth: 68 meters

Lithology:        Green mud.

Organisms:        Calcareous Benthic Foraminifers  
                   Pelecypods  
                   Ostracodes  
                   Diatoms

Ostracode Species:	Adults	Juv.	%
<u>Cytheropteron</u> sp. A	14	18	96.97
<u>Loxoconcha</u> sp. A		1	3.03
Total Ostracodes 33			

EGAL-75-KC Shipek-433

Latitude: 59° 57.5' N  
 Longitude: 141° 30.8' W  
 Water Depth: 68 meters

Lithology: Olive-green mud with some sand overlying gray mud.

Organisms: Calcareous Benthic Foraminifers  
 Pelecypods  
 Ostracodes

Ostracode Species:	Adults	Juv.,	%
<u>Cytheropteron</u> sp. A	11	26	100
Total Ostracodes 37			

REFERENCE CITED

Brouwers, E. M., 1981, Tabulation of the ostracode assemblages and associated organisms from selected bottom grab samples taken in the northeast Gulf of Alaska, F.R.S. Townsend Cromwell Cruise EGAL-75-KC, 1975: U.S. Geological Survey Open-File Report 81-1314 134 p.